

141

AGR-2002/M-221

9879 \* 36

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 6 : <b>C07D 231/20, 401/12, A01N 43/56</b>		A1	(11) International Publication Number: <b>WO 97/41106</b>
			(43) International Publication Date: 6 November 1997 (06.11.97)
(21) International Application Number: <b>PCT/JP97/01457</b>		(74) Agents: YAMAMOTO, Ryoza et al.; Torimoto Kogyo Building, 38, Kanda-Higashimatsushitacho, Chiyoda-ku, Tokyo 101 (JP).	
(22) International Filing Date: 25 April 1997 (25.04.97)			
(30) Priority Data: 8/130879 26 April 1996 (26.04.96) JP 8/227767 9 August 1996 (09.08.96) JP		(81) Designated States: BG, CA, CN, CZ, HU, KR, MX, PL, RO, RU, SI, SK, US, YU, European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).	
(71) Applicant (for all designated States except US): ISHIHARA SANGYO KAISHA LTD. [JP/JP]; 3-15, Edobori 1-chome, Nishi-ku, Osaka-shi, Osaka 550 (JP).		Published With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.	
(72) Inventors; and (75) Inventors/Applicants (for US only): MURAL, Shigeo [JP/JP]; Ishihara Sangyo Kaisha Ltd. Chuo Kenkyusho, 3-1, Nishi-shibukawa 2-chome, Kusatsu-shi, Shiga 525 (JP). KIKUGAWA, Hiroshi [JP/JP]; Ishihara Sangyo Kaisha Ltd. Chuo Kenkyusho, 3-1, Nishi-shibukawa 2-chome, Kusatsu-shi, Shiga 525 (JP). NAKAYAMA, Hitoshi [JP/JP]; Ishihara Sangyo Kaisha Ltd. Chuo Kenkyusho, 3-1, Nishi-shibukawa 2-chome, Kusatsu-shi, Shiga 525 (JP). SANO, Makiko [JP/JP]; Ishihara Sangyo Kaisha Ltd. Chuo Kenkyusho, 3-1, Nishi-shibukawa 2-chome, Kusatsu-shi, Shiga 525 (JP). ISOGAI, Akihiko [JP/JP]; Ishihara Sangyo Kaisha Ltd. Chuo Kenkyusho, 3-1, Nishi-shibukawa 2-chome, Kusatsu-shi, Shiga 525 (JP).			
(54) Title: PYRAZOLE COMPOUNDS, PROCESSES FOR THEIR PRODUCTION AND HERBICIDES CONTAINING THEM			
(57) Abstract			
<p>A pyrazole compound of formula (I) or its salt wherein R<sub>1</sub> is an alkyl group, R<sub>2</sub> is a hydrogen atom, a methyl group, -A-R<sub>3</sub>, a phenyl group which may be substituted, a pyridyl group which may be substituted, or an allyl group which is substituted by a phenyl group, A is -SO<sub>2</sub>-, -CO-, -CH(R<sub>6</sub>)- or -CH(R<sub>7</sub>)CO-, R<sub>3</sub> is an alkyl group which may be substituted, an alkenyl group which may be substituted, an alkynyl group which may be substituted, an alkoxy group which may be substituted, a cyano group, a dialkylamino group or a phenyl group which may be substituted, each of R<sub>6</sub> and R<sub>7</sub> is a hydrogen atom or an alkyl group, X is a hydrogen atom, a halogen atom, an alkyl group, a haloalkyl group, an alkoxy group, an alkylthio group, an alkylsulfinyl group, an alkylsulfonyl group, a nitro group, an alkoxy carbonyl group, -SO<sub>2</sub>N(R<sub>8</sub>)R<sub>9</sub>, -N(R<sub>10</sub>)SO<sub>2</sub>R<sub>11</sub>, -CH<sub>2</sub>S(O)<sub>q</sub>R<sub>12</sub> or -OSO<sub>2</sub>R<sub>13</sub>, each of R<sub>8</sub>, R<sub>9</sub>, R<sub>10</sub>, R<sub>11</sub>, R<sub>12</sub> and R<sub>13</sub> is an alkyl group, Z is an alkyl group, 1 is an integer of from 0 to 5, n is an integer of from 1 to 5, and q is an integer of from 0 to 2, provided that when 1 is at least 2, a plurality of Z may be the same or different, and when n is at least 2, a plurality of X may be the same or different.</p>		<p style="text-align: right;">(I)</p>	

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

DESCRIPTIONPYRAZOLE COMPOUNDS, PROCESSES FOR THEIR PRODUCTION AND  
HERBICIDES CONTAINING THEM

5

TECHNICAL FIELD

The present invention relates to novel pyrazole compounds useful as active ingredients for herbicides.

BACKGROUND ART

UK 2002375A and EP 282944A disclose pyrazole derivatives having various substituents at the 3-position of a pyrazole ring. However, the pyrazole compounds of the present invention are clearly distinguished from such derivatives in that they have a cycloalkyl group substituted at the 3-position of a pyrazole ring.

15 Further, EP 638555A discloses pyrazole glycolic acid amide derivatives having various substituents at the 3- and 4-positions of a pyrazole ring. However, the pyrazole compounds of the present invention are clearly distinguished from such derivatives in that they have a substituted benzoyl group substituted at the 4-position of a pyrazole ring.


20

DISCLOSURE OF THE INVENTION

The present inventors have conducted various studies paying attention to pyrazole compounds to find out an excellent herbicide and as a result, have accomplished the present invention. Namely, the present invention provides novel pyrazole compounds of the formula (I) or

25

their salts:



wherein  $R_1$  is an alkyl group,  $R_2$  is a hydrogen atom, a methyl group,  $-A-R_3$ , a phenyl group which may be substituted, a pyridyl group which may be substituted, or  
 10 an allyl group which is substituted by a phenyl group,  $A$  is  $-SO_2-$ ,  $-CO-$ ,  $-CH(R_6)-$  or  $-CH(R_7)CO-$ ,  $R_3$  is an alkyl group which may be substituted, an alkenyl group which may be substituted, an alkynyl group which may be substituted, an alkoxy group which may be substituted, a  
 15 cyano group, a dialkylamino group or a phenyl group which may be substituted, each of  $R_6$  and  $R_7$  is a hydrogen atom or an alkyl group,  $X$  is a hydrogen atom, a halogen atom, an alkyl group, a haloalkyl group, an alkoxy group, an alkylthio group, an alkylsulfinyl group, an alkylsulfonyl group, a nitro group, an alkoxycarbonyl group,  
 20  $-SO_2N(R_8)R_9$ ,  $-N(R_{10})SO_2R_{11}$ ,  $-CH_2S(O)qR_{12}$  or  $-OSO_2R_{13}$ , each of  $R_8$ ,  $R_9$ ,  $R_{10}$ ,  $R_{11}$ ,  $R_{12}$  and  $R_{13}$  is an alkyl group,  $Z$  is an alkyl group,  $l$  is an integer of from 0 to 5,  $n$  is an integer of from 1 to 5, and  $q$  is an integer of from 0 to  
 25 2, provided that when  $l$  is at least 2, a plurality of  $Z$  may be the same or different, and when  $n$  is at least 2, a

plurality of X may be the same or different; processes for their production; herbicides containing them; and novel intermediate compounds useful for producing them.

Now, the present invention will be described in  
5 detail with reference to the preferred embodiments.

The alkyl group or the alkyl moiety for  $R_1$  and  $R_3$  may be a  $C_{1-10}$ , preferably  $C_{1-5}$ , linear or branched alkyl group, and the alkyl group for  $R_6$  and  $R_7$  may be a  $C_{1-2}$  alkyl group. The alkyl group or the alkyl moiety for  $R_8$ ,  
10  $R_9$ ,  $R_{10}$ ,  $R_{11}$ ,  $R_{12}$ ,  $R_{13}$ , X and Z may be a  $C_{1-4}$  linear or branched alkyl group. Specific examples of such an alkyl group or moiety include methyl, ethyl, propyl, isopropyl, butyl, tert-butyl, pentyl, octyl and decyl. The alkenyl group for  $R_3$  may be a  $C_{2-10}$  linear or branched alkenyl  
15 group, such as vinyl, allyl, butadienyl or isopropenyl. The alkynyl group for  $R_3$  may be a  $C_{2-10}$  linear or branched alkynyl group, such as ethynyl, propynyl or 2-penten-4-ynyl.

The substituent for the phenyl group which may be  
20 substituted or the pyridyl group which may be substituted, for  $R_2$ , may be halogen,  $C_{1-4}$  haloalkyl or nitro. The number of substituents may be one or more, and when the number is at least 2, a plurality of such substituents may be the same or different.

25 The substituent for the alkyl which may be substituted, the alkenyl which may be substituted, the alkynyl which may be substituted, or the alkoxy group

which may be substituted, for  $R_3$ , may be halogen,  $C_{1-4}$  alkoxy,  $C_{1-6}$  alkoxycarbonyl or cyano. The number of substituents may be one or more, and if it is at least 2, a plurality of such substituents may be the same or  
5 different.

The substituent for the phenyl group which may be substituted, for  $R_3$ , may be halogen,  $C_{1-4}$  alkyl,  $C_{1-4}$  haloalkyl,  $C_{1-4}$  alkoxy- $C_{1-4}$  alkyl,  $C_{1-4}$  alkoxy, nitro or cyano. The number of substituents may be one or more,  
10 and if it is at least 2, a plurality of such substituents may be the same or different.


The halogen atom for X and the halogen as the substituent contained in  $R_2$ ,  $R_3$  and X, may be a fluorine atom, a chlorine atom, a bromine atom or an iodine atom.  
15 The number of halogen atoms as substituents, may be one or more, and if it is at least 2, a plurality of halogen atoms may be the same or different.

Among pyrazole compounds of the formula (I), a compound wherein  $R_2$  is a hydrogen atom, is capable of  
20 forming a salt. The salt may be any salt so long as it is agriculturally acceptable, and it may, for example, be an alkali metal salt such as a sodium salt or a potassium salt, an alkaline earth metal salt such as a magnesium salt or a calcium salt, or an ammonium salt such as a  
25 dimethylamine salt or a triethylamine salt.


The pyrazole compounds of the formula (I) or their salts (hereinafter referred to as the compounds of the

present invention) can be prepared in accordance with the following reactions (A) to (E) and conventional methods for producing salts.


(A) When  $R_2$  is a hydrogen atom:




(B) When  $R_2$  is a hydrogen atom:




(C) When  $R_2$  is a hydrogen atom:



(D) When  $R_2$  is a hydrogen atom:




(E) When  $R_2$  is other than a hydrogen atom:






Among the compounds of the present invention, those having certain predetermined substituents can be prepared in accordance with the following reactions (F) to (G) and conventional methods for preparing salts.

- 5 (F) When  $R_2$  is a hydrogen atom, and  $(X)_n$  contains at least one alkylsulfinyl or alkylsulfonyl group:



8

(G) When  $R_2$  is other than a hydrogen atom, and  $(X)_n$  contains at least one alkylsulfinyl or alkylsulfonyl group:



Now, the above reaction (A) will be described. In the reaction (A),  $R_1$ ,  $X$ ,  $Z$ ,  $l$  and  $n$  are as defined above, and  $Y$  is a halogen atom.

The condensation reaction in the reaction (A) can be carried out, if necessary, in the presence of a base. As such a base, one or more members may be suitably selected for use from carbonates such as potassium carbonate and sodium carbonate; hydrogencarbonates such as potassium hydrogencarbonate and sodium hydrogencarbonate; metal hydrides such as potassium hydride and sodium hydride;

amines such as monomethylamine, dimethylamine and triethylamine; and pyridines such as pyridine and 4-dimethylaminopyridine.

Further, the condensation reaction in the reaction  
5 (A) can be carried out, if necessary, in the presence of a solvent. As such a solvent, any solvent may be used so long as it is a solvent inert to the reaction, and one or more members may be suitably selected for use from aromatic hydrocarbons such as benzene, toluene, xylene  
10 and chlorobenzene; cyclic or noncyclic aliphatic hydrocarbons such as carbon tetrachloride, methylene chloride, chloroform, dichloromethane, dichloroethane, trichloroethane, hexane and cyclohexane; ethers such as dioxane, tetrahydrofuran and diethyl ether; esters such  
15 as methyl acetate and ethyl acetate; polar aprotic solvents such as dimethylsulfoxide, sulfolane, dimethylacetamide, dimethylformamide, N-methylpyrrolidone, pyridine and hexamethylphosphoric triamide; nitriles such as acetonitrile, propionitrile  
20 and acrylonitrile; ketones such as acetone and methyl ethyl ketone; and water.

Further, the condensation reaction in the reaction  
(A) can be carried out, if necessary, in the presence of a phase transfer catalyst. As such a phase transfer  
25 catalyst, one or more members may be suitably selected for use from e.g. benzyltriethylammonium chloride, benzyltriethylammonium bromide, tetraethylammonium

chloride and tetraethylammonium bromide.

The reaction temperature of the condensation reaction in the reaction (A) is usually from 0 to 250°C, preferably from 15 to 150°C, and the reaction time is  
5 usually from 0.1 to 48 hours, preferably from 0.1 to 24 hours.

The compound of the formula (IV) which can be produced by the condensation reaction in this reaction (A), is a novel intermediate compound useful for  
10 producing the compounds of the present invention.

The rearrangement reaction in the reaction (A) comprises the following two steps i.e. (1) a rearrangement reaction step and (2) a pH adjusting reaction step. The rearrangement reaction step is  
15 carried out usually in the presence of a base. As such a base, one or more members may be suitably selected for use from carbonates such as potassium carbonate and sodium carbonate; and calcium hydroxide. The base is used usually in an amount of from 0.5 to 5 mols per mol  
20 of the compound of the formula (IV).

Further, the rearrangement reaction step of the rearrangement reaction in the reaction (A) can be carried out, if necessary, in the presence of a solvent. As such a solvent, any solvent may be used so long as it is a  
25 solvent inert to the reaction, and one or more members may suitably be selected for use from aromatic hydrocarbons such as benzene, toluene, xylene and

chlorobenzene; ethers such as dioxane, tetrahydrofuran and diethyl ether; and polar aprotic solvents such as dimethylsulfoxide, sulfolane, dimethylacetamide, dimethylformamide, N-methylpyrrolidone, pyridine and  
5 hexamethylphosphoric triamide.

The rearrangement reaction step of the rearrangement reaction in the reaction (A) is preferably carried out under an azeotropic dehydrating condition, whereby the rearrangement reaction will effectively proceed. This is  
10 one of preferred embodiments of the present invention. By the rearrangement reaction step, a salt of the compound of the formula (I) is produced, and a method for producing such a salt is also one of embodiments of the present invention. Further, a compound of the above-  
15 mentioned formula (I-2) can be produced by reacting a salt of the compound of the above formula (I) or a reaction mixture containing such a salt, obtained by this rearrangement reaction step, with a compound of the above formula (VII), under the reaction conditions for the  
20 reaction (D) which will be described hereinafter. This is also one of embodiments of the present invention.

The reaction temperature in the rearrangement reaction step is usually from 50 to 250°C, preferably from 50 to 150°C, and the reaction time is usually from  
25 0.1 to 48 hours, preferably from 0.5 to 24 hours.

The pH adjusting reaction step of the rearrangement reaction in the reaction (A) is a reaction to adjust the

pH value to at most 7, which is carried out usually in the presence of an acidic substance and water. As such an acidic substance, one or more members may suitably be selected for use from inorganic acids such as  
5 hydrochloric acid and sulfuric acid; and organic acids such as acetic acid.

The pH adjusting reaction step of the rearrangement reaction in the reaction (A) can be carried out, if necessary, in the presence of a solvent. As such a  
10 solvent, any solvent may be used so long as it is a solvent inert to the reaction. For example, one or more members may be suitably selected for use from those mentioned in the description of the rearrangement reaction step as the preceding step.

15 The pH adjusting reaction step of the rearrangement reaction in the reaction (A) may be carried out after isolating the reaction product obtained by the rearrangement reaction step as the preceding step, in accordance with a conventional method, or may be carried  
20 out in one pot by using the reaction mixture obtained by the rearrangement reaction step, as it is. When it is carried out in one pot, it is carried out by adding and reacting an acidic substance and water to the reaction mixture obtained by the rearrangement reaction step as  
25 the preceding step.

The reaction temperature for the pH adjusting reaction step is usually from 0 to 100°C, preferably from

0 to 60°C.

Now, the above-mentioned reaction (B) will be described. In the reaction (B),  $R_1$ , Z, l, n and (II) are as defined above, and X' is a hydrogen atom, a halogen atom, an alkyl group, a haloalkyl group, an alkoxy group, 5 an alkylthio group, an alkylsulfinyl group or an alkylsulfonyl group, provided that when n is at least 2, a plurality of X' may be the same or different.

The condensation reaction in the reaction (B) is 10 carried out usually in the presence of a Lewis acid. As such a Lewis acid, one or more members may suitably be selected for use from e.g. dry aluminum chloride and dry aluminum bromide.

Further, the condensation reaction in the reaction 15 (B) can be carried out, if necessary, in the presence of a solvent. As such a solvent, any solvent may be used so long as it is a solvent inert to the reaction, and one or more members may suitably be selected for use from halogenated aliphatic hydrocarbons such as carbon 20 tetrachloride, methylene chloride, chloroform, dichloromethane, and dichloroethane.

The reaction temperature for the condensation reaction in the reaction (B) is usually from 0 to 80°C, and the reaction time is usually from 0.1 to 24 hours, 25 preferably from 0.1 to 10 hours.

The hydrolytic reaction in the reaction (B) is carried out usually in the presence of an acidic

substance. As such an acidic substance, one or more members may suitably be selected for use from e.g. inorganic acids such as hydrochloric acid and sulfuric acid.

5       The hydrolytic reaction in the reaction (B) can be carried out, if necessary, in the presence of a solvent. As such a solvent, any solvent may be used so long as it is a solvent inert to the reaction, and one or more members may suitably be selected for use among those  
10 exemplified in the description of the condensation reaction as the preceding reaction.

      The hydrolytic reaction in the reaction (B) may be carried out after isolating the reaction product obtained by the condensation reaction as the preceding reaction,  
15 in accordance with a conventional method, or may be carried out in one pot using the reaction mixture obtained by the condensation reaction as it is. In the case where it is carried out in one pot, post treatment such as removal of the Lewis acid may be applied, if  
20 necessary, to the reaction mixture obtained by the condensation reaction as the preceding reaction, and the acidic substance and water are added thereto to carry out the reaction.

      The reaction temperature for the hydrolytic reaction  
25 in the reaction (B) is usually from 20 to 100°C, and the reaction time is usually from 0.1 to 24 hours, preferably from 0.1 to 10 hours.



Now, the above-mentioned reaction (C) will be described. In the reaction (C), X, n, (II) and (I-1) are as defined above.

The condensation reaction in the reaction (C) is  
5 carried out usually in the presence of a condensing agent and a solvent. As such a condensing agent, N,N'-dicyclohexylcarbodiimide may, for example, be mentioned, and as such a solvent, any solvent may be used so long as it is a solvent inert to the reaction, and one or more  
10 members may suitably be selected for use among alcohols such as tert-butyl alcohol and tert-amyl alcohol.

The condensation reaction in the reaction (C) can be carried out, if necessary, in the presence of a base. As such a base, one or more members may suitably be selected  
15 for use from e.g. carbonates such as potassium carbonate and sodium carbonate.

The reaction temperature for the condensation reaction in the reaction (C) is usually from 50 to 100°C, and the reaction time is usually from 0.1 to 24 hours,  
20 preferably from 0.5 to 20 hours.

Now, the above-mentioned reaction (D) will be described. In the reaction (D), X, n, (II) and (I-1) are as defined above, and T is a chlorine atom, a bromine atom or an iodine atom.

25 The reaction (D) is carried out usually in the presence of a base and a metal catalyst. As a base, one or more members may suitably be selected for use from

e.g. alkali metals such as sodium and potassium; alkali metal alkoxides such as sodium methylate, sodium ethylate and potassium tert-butoxide; carbonates such as potassium carbonate and sodium carbonate; hydrogencarbonates such  
5 as potassium hydrogencarbonate and sodium hydrogencarbonate; metal hydroxides such as potassium hydroxide and sodium hydroxide; metal hydrides such as potassium hydride and sodium hydride; amines such as monomethylamine, dimethylamine and triethylamine;  
10 pyridines such as pyridine and 4-dimethylaminopyridine; and N,N-dimethylaniline. As the metal catalyst, a transition metal such as palladium, rhodium, ruthenium or platinum, may be mentioned. The ligand used against the metal of the metal catalyst is not particularly limited,  
15 but an organophosphine compound such as triphenylphosphine or tri-n-butylphosphine is preferred.

The reaction (D) may be carried out, if necessary, in the presence of a solvent. As such a solvent, any solvent may be used so long as it is a solvent inert to  
20 the reaction. For example, one or more members may suitably be selected for use among aromatic hydrocarbons such as benzene, toluene, xylene and chlorobenzene; cyclic or noncyclic aliphatic hydrocarbons such as carbon tetrachloride, methylene chloride, chloroform,  
25 dichloromethane, dichloroethane, trichloroethane, hexane and cyclohexane; ethers such as dioxane, tetrahydrofuran and diethyl ether; esters such as methyl acetate and

ethyl acetate; polar aprotic solvents such as dimethylsulfoxide, sulfolane, dimethylacetamide, dimethylformamide, N-methylpyrrolidone and pyridine; nitriles such as acetonitrile, propionitrile and  
5 acrylonitrile; ketones such as acetone and methyl ethyl ketone; amines such as monomethylamine, dimethylamine and triethylamine; alcohols such as methanol, ethanol, propanol, and tert-butanol; organic acids such as acetic acid and propionic acid; aqueous ammonia; and water.

10 The reaction temperature for the reaction (D) is usually from 30 to 300°C, preferably from 50 to 200°C, and the reaction time is usually from 0.1 to 48 hours, preferably from 1 to 24 hours.

Now, the above-mentioned reaction (E) will be  
15 described. In the reaction (E),  $R_1$ , X, Y, Z, l, n and (I-1) are as defined above, and  $R_2'$  is a methyl group, -A- $R_3$ , a phenyl group which may be substituted, a pyridyl group which may be substituted or an allyl group which is substituted by a phenyl group (wherein A and  $R_3$  are as  
20 defined above).

The condensation reaction in the reaction (E) may be carried out, if necessary, in the presence of a base. As such a base, one or more members may suitably be selected for use from carbonates such as potassium carbonate and  
25 sodium carbonate; hydrogencarbonates such as potassium hydrogencarbonate and sodium hydrogencarbonate; metal hydroxides such as potassium hydroxide and sodium

hydroxide; metal hydrides such as potassium hydride and sodium hydride; amines such as monomethylamine, dimethylamine and triethylamine; and pyridines such as pyridine and 4-dimethylaminopyridine.

5       The condensation reaction in the reaction (E) may be carried out, if necessary, in the presence of a solvent. As such a solvent, any solvent may be used so long as it is inert to the reaction. For example, one or more members may suitably be selected for use from aromatic  
10   hydrocarbons such as benzene, toluene, xylene and chlorobenzene; cyclic or noncyclic aliphatic hydrocarbons such as carbon tetrachloride, methylene chloride, chloroform, dichloromethane, dichloroethane, trichloroethane, hexane and cyclohexane; ethers such as  
15   dioxane, tetrahydrofuran and diethyl ether; esters such as methyl acetate and ethyl acetate; polar aprotic solvents such as dimethylsulfoxide, sulfolane, dimethylacetamide, dimethylformamide, N-methylpyrrolidone, pyridine and hexamethylphosphoric  
20   triamide; nitriles such as acetonitrile, propionitrile and acrylonitrile; ketones such as acetone and methyl ethyl ketone; and water.

      The condensation reaction in the reaction (E) may be carried out, if necessary, in the presence of a phase  
25   transfer catalyst and/or potassium iodide. As such a phase transfer catalyst, one or more members may suitably be selected for use among those mentioned for the

condensation reaction in the above-mentioned reaction (A).

The reaction temperature for the condensation reaction in the reaction (E) is usually from 0 to 200°C, preferably from 15 to 150°C, and the reaction time is usually from 0.1 to 48 hours, preferably from 0.1 to 24 hours.

Now, the above-mentioned reaction (F) will be described. In the reaction (F),  $R_1$ , X, Z, l and n are as defined above,  $R_5$  is an alkyl group, preferably a  $C_{1-4}$  alkyl group, X" is a hydrogen atom, a halogen atom, an alkyl group, a haloalkyl group, an alkoxy group, an alkylsulfinyl group, an alkylsulfonyl group, a nitro group, an alkoxy carbonyl group,  $-SO_2N(R_8)R_9$ ,  $-N(R_{10})SO_2R_{11}$ ,  $-CH_2S(O)_{q'}R_{12}$  or  $-OSO_2R_{13}$ , wherein  $R_8$ ,  $R_9$ ,  $R_{10}$ ,  $R_{11}$ ,  $R_{12}$  and  $R_{13}$  are as defined above, m is 1 or 2, and  $q'$  is 1 or 2. In the reaction (F), the oxidation reaction for producing (IV-2) from (IV-1) and the oxidation reaction for producing (I-4) from (I-5) (hereinafter referred to simply as the oxidation reaction) are carried out usually in the presence of an oxidizing agent and a solvent. As such an oxidizing agent, one or more members may suitably be selected for use from e.g. m-chloroperbenzoic acid and hydrogen peroxide. As the solvent, any solvent may be used so long as it is a solvent inert to the reaction. For example, one or more members may suitably be selected for

use among those mentioned for the condensation reaction in the above-mentioned reaction (B).

The reaction temperature for the oxidation reaction in the reaction (F) is usually from 0 to 80°C, and the  
5 reaction time is usually from 0.1 to 48 hours, preferably from 0.1 to 24 hours.


The rearrangement reaction in the reaction (F) can be carried out in accordance with the rearrangement reaction in the above-mentioned reaction (A).

10 Now, the above-mentioned reaction (G) will be described. In the reaction (G),  $R_1$ ,  $R_2'$ ,  $R_5$ ,  $X$ ,  $X''$ ,  $Z$ ,  $l$ ,  $m$  and  $n$  are as defined above.

The oxidation reaction in the reaction (G) can be carried out in accordance with the oxidation reaction in  
15 the above-mentioned reaction (F).

The compound of the formula (II) in the above reactions (A), (B), (C) and (D) is a novel intermediate compound which is useful for producing the compounds of the present invention and may be produced, for example,  
20 by a method such as the reaction (H).

(H)



Now, the reaction (H) will be described. In the reaction (H),  $R_1$ ,  $Y$ ,  $Z$  and  $l$  are as defined above, and  $R_4$  is a  $C_{1-6}$  alkyl group.

In the reaction (H), the cyclization reaction for producing (II) from (VIII) and the cyclization reaction for producing (IX) from (VIII) (hereinafter referred to simply as the cyclization reaction) may be carried out, if necessary, in the presence of a solvent. As such a solvent, any solvent may be used so long as it is a solvent inert to the reaction. For example, one or more members may suitably be selected for use from aromatic

hydrocarbons such as benzene, toluene, xylene and chlorobenzene; cyclic or noncyclic aliphatic hydrocarbons such as carbon tetrachloride, methylene chloride, chloroform, dichloromethane, dichloroethane, trichloroethane, hexane and cyclohexane; ethers such as dioxane, tetrahydrofuran and diethyl ether; polar aprotic solvents such as dimethylsulfoxide, sulfolane, dimethylacetamide, dimethylformamide, N-methylpyrrolidone pyridine and hexamethylphosphoric triamide; nitriles such as acetonitrile, propionitrile and acrylonitrile; and water.

The cyclization reaction in the reaction (H) may be carried out, if necessary, under an azeotropic dehydration condition.

The reaction temperature for the cyclization reaction in the reaction (H) is usually from 0 to 200°C, preferably from 20 to 150°C, and the reaction time is usually from 0.1 to 48 hours, preferably from 0.1 to 24 hours.

The condensation reaction in the reaction (H) is carried out usually in the presence of a base and a solvent. As the base, one or more members may suitably be selected for use from carbonates such as potassium carbonate and sodium carbonate; and metal hydrides such as potassium hydride and sodium hydride. Particularly preferred is potassium carbonate.

As the solvent, any solvent may be used so long as it






is a solvent inert to the reaction. For example, one or more members may suitably be selected for use from ethers such as dioxane, tetrahydrofuran and diethyl ether; and polar aprotic solvents such as dimethylsulfoxide, 5 sulfolane, dimethylacetamide, dimethylformamide, N-methylpyrrolidone, pyridine and hexamethylphosphoric triamide. Particularly preferred is hexamethylphosphoric triamide.

The reaction temperature for the condensation 10 reaction in the reaction (H) is usually from -20 to +150°C, preferably from -15 to +60°C, and the reaction time is usually from 0.1 to 24 hours, preferably from 0.1 to 10 hours.

The compound of the formula (IX) which can be 15 prepared by the cyclization reaction in this reaction (H), is a novel intermediate compound which is useful for producing the compounds of the present invention.






The compounds of the present invention and the intermediate compounds useful for the production thereof, 20 have the following isomers. Such various isomers (the respective isomers and mixtures of such isomers) are within the scope of the present invention.

(1) Among the compounds of the present invention represented by the above formula (I), compounds wherein 25  $R_2$  is a hydrogen atom, and intermediate compounds represented by the above formulas (II) and (IX), have the following tautomers, respectively.



wherein  $R_1$ ,  $X$ ,  $Z$ ,  $l$  and  $n$  are as defined above.

(2) Among the compounds of the present invention represented by the above formula (I) and the intermediate compounds represented by the above formulas (II), (IV), (VIII) and (IX), compounds wherein 1 is at least 1, have optical isomers. Some examples will be given below, but it should be understood that the optical isomers in the present invention are not limited to such specific examples.



wherein  $R_1$ ,  $R_2$ ,  $R_4$ ,  $X$ ,  $Z$  and  $n$  are as defined above.

In the specification of this application, such optical isomers are meant for a mixture of isomers (racemic modification) unless otherwise specified.

(3) Among compounds of the present invention represented by the above formula (I), compounds wherein  $R_2$  is a  $-A-R_3$ , and  $R_3$  is an alkenyl group which may be substituted, have geometrical isomers (E-isomer and Z-isomer).

The compound of the present invention exhibits excellent herbicidal effects when used as an active ingredient of a herbicide. It finds a wide range of applications to crop lands such as paddy fields, upland farms, orchards and mulberry fields, and non-crop lands such as forests, farm roads, playgrounds, and factory sites. The application method may suitably be selected from soil treatment application and foliar application.

The herbicidal composition containing the compound of the present invention is capable of controlling noxious weeds including grasses (or gramineae) such as barnyardgrass (Echinochloa crus-galli L.), crabgrass (Digitaria sanguinalis L.), greenfoxtail (Setaria viridis L.), goosegrass (Eleusine indica L.), wild oat (Avena fatua L.), johnsongrass (Sorghum halepense L.), quackgrass (Agropyron repens L.), alexandergrass (Brachiaria plantaginea), paragrass (Panicum purpurascens), sprangletop (Leptochloa chinensis) and red sprangletop (Leptochloa panicea); sedges (or Cyperaceae) such as rice flatsedge (Cyperus iria L.), purple nutsedge (Cyperus rotundus L.), japanese bulrush (Scirpus juncoides), flatsedge (Cyperus serotinus), small-flower

- umbrellaplant (Cyperus difformis), slender spikerush (Eleocharis acicularis), and water chestnut (Eleocharis kuroquwai); alismataceae such as japanese ribbon wapato (Sagittaria pygmaea), arrow-head (Sagittaria trifolia) and narrowleaf waterplantain (Alisma canaliculatum); pontederiaceae such as monochoria (Monochoria vaginalis) and monochoria species (Monochoria korsakowii); scrophulariaceae such as false pimpernel (Lindernia pyxidaria) and abunome (Dopatrium junceum); lythraceae such as toothcup (Rotala indica) and red stem (Ammannia multiflora); and broadleaves such as velvetleaf (Abutilon theophrasti MEDIC.), tall morningglory (Ipomoea purpurea L.), common lambsquarters (Chenopodium album L.), prickly sida (Sida spinosa L.), common purslane (Portulaca oleracea L.), slender amaranth (Amaranthus viridis L.), redroot pigweed (Amaranthus retroflexus L.), sicklepod (Cassia obtusifolia L.), black nightshade (Solanum nigrum L.), pale smartweed (Polygonum lapathifolium L.), common chickweed (Stellaria media L.), common cocklebur (Xanthium strumarium L.), henbit (Lamium amplexicaule L.) and flexuosa WITH.), flexuous bittercress (Cardamine threeseeded copperleaf (Acalypha australis L.).
- Accordingly, it is useful for controlling noxious weeds non-selectively or selectively in the cultivation of a crop plant such as corn (Zea mays L.), soybean (Glycine max Merr.), cotton (Gossypium spp.), wheat (Triticum spp.), rice (Oryza sativa L.), barley (Hordeum

vulgare L.), oat (Avena sativa L.), sorgo (Sorghum  
bicolor Moench), rape (Brassica napus L.), sunflower  
(Helianthus annuus L.), sugar beet (Beta vulgaris L.),  
sugar cane (Saccharum officinarum L.), japanese lawngress  
5 (Zoysia japonica stend), peanut (Arachis hypogaea L.) or  
flax (Linum usitatissimum L.) The compound of the  
present invention is particularly effective for  
selectively controlling noxious weeds in the cultivation  
of corn, wheat or rice, especially in the cultivation of  
10 corn.

The herbicidal composition containing the compound of  
the present invention is usually formulated by mixing the  
compound with various agricultural adjuvants and used in  
the form of a formulation such as a dust, granules,  
15 water-dispersible granules, a wettable powder, an  
emulsifiable concentrate, a water-based suspension  
concentrate, an oil-based suspension concentrate, water  
soluble granules (or powder), tablets or capsules.  
However, so long as it is suitable for the purpose of the  
20 present invention, it may be formulated into any type of  
formulation which is commonly used in this field.

Such agricultural adjuvants include solid carriers  
such as diatomaceous earth, slaked lime, calcium  
carbonate, talc, white carbon, kaoline, bentonite, a  
25 mixture of kaolinite and sericite, clay, sodium  
carbonate, sodium bicarbonate, mirabilite, zeolite and  
starch; solvents such as water, toluene, xylene, solvent

naphtha, dioxane, acetone, isophorone, methyl isobutyl ketone, chlorobenzene, cyclohexane, dimethylsulfoxide, dimethylformamide, N-methyl-2-pyrrolidone, and alcohol; anionic surfactants and spreaders such as a salt of fatty acid, a benzoate, an alkylsulfosuccinate, a dialkylsulfosuccinate, a polycarboxylate, a salt of alkylsulfuric acid ester, an alkyl sulfate, an alkylaryl sulfate, an alkyl diglycol ether sulfate, a salt of alcohol sulfuric acid ester, an alkyl sulfonate, an alkylaryl sulfonate, an aryl sulfonate, a lignin sulfonate, an alkyl diphenyl ether disulfonate, a polystyrene sulfonate, a salt of alkylphosphoric acid ester, an alkylaryl phosphate, a styrylaryl phosphate, a salt of polyoxyethylene alkyl ether sulfuric acid ester, a polyoxyethylene alkylaryl ether sulfate, a salt of polyoxyethylene alkyl ether sulfuric acid ester, a polyoxyethylene alkylaryl ether sulfuric acid ester, a polyoxyethylene alkyl ether phosphate, a salt of polyoxyethylene alkyl aryl phosphoric acid ester, and a salt of a condensate of naphthalene sulfonate with formalin; nonionic surfactants and spreaders such as a sorbitan fatty acid ester, a glycerin fatty acid ester, a fatty acid polyglyceride, a fatty acid alcohol polyglycol ether, acetylene glycol, acetylene alcohol, an oxyalkylene block polymer, a polyoxyethylene alkyl ether, a polyoxyethylene alkylaryl ether, a polyoxyethylene styrylaryl ether, a polyoxyethylene glycol alkyl ether, a

polyoxyethylene fatty acid ester, a polyoxyethylene sorbitan fatty acid ester, a polyoxyethylene glycerin fatty acid ester, a polyoxyethylene hydrogenated castor oil, and a polyoxypropylene fatty acid ester; and  
5 vegetable and mineral oils such as olive oil, kapok oil, castor oil, palm oil, camellia oil, coconut oil, sesame oil, corn oil, rice bran oil, peanut oil, cottonseed oil, soybean oil, rapeseed oil, linseed oil, tung oil, and liquid paraffins. Such adjuvants may be selected for use  
10 among those known in this field, so long as the purpose of the present invention can thereby be accomplished. Further, various additives which are commonly used, such as a filler, a thickener, an anti-settling agent, an anti-freezing agent, a dispersion stabilizer, a  
15 phytotoxicity reducing agent, and an anti-mold agent, may also be employed.

The weight ratio of the compound of the present invention to the various agricultural adjuvants is usually from 0.1 :99.9 to 95:5, preferably from 0.2:99.8  
20 to 85:15.

The dose of the herbicidal composition of the present invention can not generally be defined, since it may vary depending upon the weather condition, the soil condition, the type of the formulation, the types of the weeds to be  
25 controlled, the season for the application, etc. However, it is usually applied so that the compound of the present invention would be applied in an amount of



from 0.5 to 5000 g/ha, preferably from 1 to 1000 g/ha, more preferably from 5 to 500 g/ha. The present invention covers such a method for controlling noxious weeds by application of such a herbicidal composition.

5       The herbicidal compositions of the present invention may be used in admixture with or in combination with other agricultural chemicals, fertilizers or phytotoxicity-reducing agents. In such a case, they may exhibit even better effects or activities. As other  
10       agricultural chemicals, herbicides, fungicides, antibiotics, plant hormones or insecticides may, for example, be mentioned. Especially with a mixed herbicidal composition having the compound of the present invention used in admixture with or in combination with  
15       one or more active ingredients of other herbicides, it is possible to improve the herbicidal activities, the season for the application and the range of applicable weed types. Further, the compound of the present invention and an active ingredient of other herbicide may be  
20       separately formulated, so that they may be mixed for use at the time of application, or both may be formulated together. The present invention covers such mixed herbicidal compositions.

      The blend ratio of the compounds of the present  
25       invention with the active ingredients of other herbicides can not generally be defined, since it varies depending upon the weather condition, the soil condition, the type

of the formulation, the season for the application, the manner of the application, etc. However, one active ingredient of other herbicide may be incorporated usually in an amount of from 0.001 to 10000 parts by weight, preferably from 0.01 to 1000 parts by weight, per part by weight of the compound of the present invention. Further, the total dose of all of the active ingredients is usually from 0.1 to 10000 g/ha, preferably from 0.2 to 5000 g/ha. The present invention covers a method for controlling noxious weeds by application of such herbicidal compositions.

As the active ingredients of other herbicides, the following (common names) may be mentioned.

(1) Those which are believed to exhibit herbicidal effects by disturbing auxin activities of plants, including a phenoxy acetic acid type such as 2,4-D, MCPA, MCPB or naproanilide, an aromatic carboxylic acid type such as 2,3,6-TBA, dicamba, picloram or clopyralid, and others such as benazolin, quinclorac, quinmerac or diflufenzopyr.

(2) Those which are believed to exhibit herbicidal effects by inhibiting photosynthesis of plants, including a urea type such as diuron, linuron, isoproturon or metobenzuron, triazine type such as simazine, atrazine, atratone, simetryn, prometryn, dimethametryn, metribuzin, terbuthylazine, cyanazine or ametryn, an uracil type such as bromacil or lenacil, an anilide type such as propanil

or cypromid, a carbamate type such as swep or phenmedipham, a hydroxybenzonitrile type such as bromoxynil, bromoxynil-octanoate or ioxynil, and others such as pyridate or bentazon.

5 (3) A quaternary ammonium salt type such as paraquat or diguat, which is believed to be converted to free radicals by itself to form active oxygen in the plant body and thus to exhibit quick herbicidal effects.

(4) Those which are believed to exhibit herbicidal effects by inhibiting chlorophyllbiosynthesis of plants and abnormally accumulating a photosensitizing peroxide substance in the plant body, including a diphenyl ether type such as nitrofen, chlomethoxyfen, bifenox, acifluorfen-sodium, fomesafen or oxyfluorfen, a cyclic imide type such as chlorphthalim, flumioxadine, flumiclorac-pentyl, methyl [2-chloro-4-fluoro-5-(5,6,7,8-tetrahydro-3-oxo-1H,3H-[1,3,4]thiadiazolo[3,4-a]pyridazin-1-ylideneamino)phenylthio] acetate (compound disclosed at page 60 of proceedings of 19th Meeting of Pesticide Science Society of Japan), and others such as oxadiation, sulfentrazone, carfentrazone-ethyl, thidiazimin, ethyl 2-chloro-5-(4-chloro-5-difluoromethoxyl-1-methylpyrazol-3-yl)-4-fluorophenoxyacetate (compound disclosed at pages 70-71 of proceedings of 21th Meeting of Pesticide Science Society of Japan).

(5) Those which are believed to exhibit herbicidal

- effects characterized by whitening activities by inhibiting chromogenesis of plants such as carotenoids, including a pyridazinone type such as norflurazon or metflurazon, a pyrazole type such as pyrazolate, 5 pyrazoxyfen or benzofenap, and others such as fluridone, flurtamone, diflufenican, methoxyphenone, clomazone, sulcotrione, 2-(2'-nitro-4'-methylsulfonyl-benzoyl)-1,3-cyclohexanedione (compound disclosed in US Patent 5,506,195), isoxaflutole or difenzoquat.
- 10 (6) Those which exhibit herbicidal effects specifically to gramineous plants, including an aryloxyphenoxypropionic acid type such as diclofop-methyl, pyriphenop-sodium, fluazifop-butyl, haloxyfop-methyl, quizalofop-ethyl or cyhalofop-butyl, and a 15 cyclohexanedione type such as alloxydim-sodium, clethodim, sethoxydim or tralkoxydim.
- (7) Those which are believed to exhibit herbicidal effects by inhibiting an amino acid biosynthesis of plants, including a sulfonylurea type such as 20 chlorimuron-ethyl, sulfometuron-methyl, primisulfuron-methyl, bensulfuron-methyl, chlorsulfuron, metsulfuron-methyl, cinosulfuron, pyrazosulfuron-ethyl, azimsulfuron, flazasulfuron, rimusulfuron, nicosulfuron, imazosulfuron, cyclosulfamuron, prosulfuron, flupyrsulfuron, 25 trisulfuron-methyl, halosulfuron-methyl or thifensulfuron-methyl, a triazolopyrimidinesulfoneamide type such as flumetsulam or metosulam, an imidazolinone

type such as imazapyr, imazethapyr, imazaquin, imazamox or imazameth or imazamethabenz, a pyrimidinylsalicylic acid type such as pyriithiobac-sodium, bispyribac-sodium or pyriminobac-methyl, and others such as glyphosate-  
5 ammonium, glyphosate-isopropylamine, glufosinate-ammonium or bialaphos.

(8) Those which are believed to exhibit herbicidal effects by inhibiting cell mitoses of plants, including a dinitroaniline type such as trifluralin, oryzalin,  
10 nitralin or pendimethalin, an organic phosphorus type such as amiprofos-methyl, butamifos, anilofos or piperophos, a phenylcarbamate type such as chlorpropham or barban, a cumylamine type such as daimuron, cumyluron or bromobutide, and others such as asulam or dithiopyr.

15 (9) Those which are believed to exhibit herbicidal effects by inhibiting protein biosynthesis or lipid biosynthesis of plants, including a thiocarbamate type such as EPTC, butylate, molinate, dimepiperate, esprocarb, thiobencarb or pyributicarb, or  
20 chloroacetamide type such as alachlor, butachlor, pretilachlor, metolachlor, thenylchlor, dimethenamid, acetochlor or propachlor, and other compounds such as a ethobenzanide, mefenacet, thiafluamide, tridiphane, cafenstrole, 4-(2-chlorophenyl)-N-cyclohexyl-4,5-dihydro-  
25 N-ethyl-5-oxo-1H-tetrazol-1-carboxamide (compound disclosed in JP-A-6-306061), oxaziclomefon, or 2-ethyl-2-[2-(3-chlorophenyl)-2,3-epoxypropyl]-indan-1,3-dione

(compound disclosed in JP-A-2-304043).

As is evident from Test Examples 1 and 2 given hereinafter, the compound of the present invention include those which show selectivity for effectively  
5 controlling weeds, while showing safety to crop plants such as rice, wheat and corn. When the compound of the present invention is to be used in the cultivation of such crop plants, synergistic effects may be obtained by using it in admixture with or in combination with one or  
10 more of the following compounds among the above-mentioned active compounds of other herbicides.

In the cultivation of rice:

2,4-D, MCPA, MCPB, naproanilide, quinclorac, simetryn, prometryn, dimethametryn, propanil, swep,  
15 bentazon, nitrofen, chlomethoxyfen, bifenox, oxadiazon, pyrazolate, pyrazoxyfen, benzofenap, methoxyphenone, cyhalofop-butyl, bensulfuron-methyl, cinosulfuron, pyrazosulfuron-ethyl, azimsulfuron, imazosulfuron, cyclosulfamuron, bispyribac-sodium salt, pyriminobac-  
20 methyl, anilofos, piperophos, daimuron, cumyluron, bromobutide, dithiopyr, molinate, dimepiperate, esprocarb, thiobencarb, pyributicarb, thenylchlor, pretilachlor, butachlor, ethobenzanide, mefenacet, cafenstrole, 4-(2-chlorophenyl)-N-cyclohexyl-4,5-dihydro-  
25 N-ethyl-5-oxo-1H-tetrazole-1-carboxamide, oxaziclomefon, and 2-ethyl-2-[2-(3-chlorophenyl)-2,3-epoxypropyl]-indane-1,3-dione.

In the cultivation of corn:

2,4-D, MCPA, dicamba, clopyralid, benazolin,  
diflufenzopyr, diuron, linuron, metobenzuron, simazine,  
atrazine, atratone, metribuzin, terbuthylazine,  
5 cyanazine, ametryn, cypromid, bromoxynil, bromoxynil-  
octanoate, pyridate, bentazon, paraquat, oxyfluorfen,  
flumiclorac-pentyl, methyl [2-chloro-4-fluoro-5-(5,6,7,8-  
tetrahydro-3-oxo-1H,3H-[1,3,4]thiadiazolo[3,4-  
a]pyridazin-1-ylideneamino)phenylthio] acetate,  
10 fluridone, sulcotrione, 2-(2'-nitro-4'-  
methylsulfonylbenzoyl)-1,3-cyclohexanedione,  
isoxaflutole, carfentrazone ethyl, primisulfuron methyl,  
rimusulfuron, nicosulfuron, prosulfuron, halosulfuron-  
methyl, thifensulfuron-methyl, flumetsulam, metosulam,  
15 imazethapyr, glyphosate-ammonium salt, glyphosate-  
isopropyl amine salt, glufosinate-ammonium salt,  
trifluralin, pendimethalin, EPTC, butylate, alachlor,  
metolachlor, acetochlor, propachlor, dimethenamid and  
tridiphane.

20 In the cultivation on wheat:


MCPB, quinmerac, linuron, isoproturon, prometryn,  
bromoxynil, bromoxynil-octanoate, pyridate, bifenox,  
carfentrazone-ethyl, thidiazimin, ethyl 2-chloro-5-(4-  
chloro-5-difluoromethoxyl-1-methylpyrazol-3-yl)-4-  
25 fluorophenoxy acetate, flurtamone, diflufenican,  
sulcotrione, diclofop-methyl, tralkoxydim, chlorsulfuron,  
metsulfuron-methyl, prosulfuron, halosulfuron-methyl,

flumetsulam, metosulam, pendimethalin, barban and imazamethabenz.

Now, preferred embodiments of the present invention will be described.

5 (1) The pyrazole compound of the above formula (I) or its salt.

(2) The pyrazole compound or its salt according to Item 1, wherein the formula (I) is represented by the formula (I'):



wherein  $R_1$  is an alkyl group,  $R_2$  is a hydrogen atom or  $-A-R_3$ ,  $A$  is  $-\text{SO}_2-$ ,  $-\text{CO}-$ ,  $-\text{CH}_2-$  or  $-\text{CH}_2\text{CO}-$ ,  $R_3$  is an alkyl group which may be substituted, an alkenyl group which may be substituted, an alkynyl group which may be substituted, a cyano group or a phenyl group which may be substituted, each of  $X^1$ ,  $X^2$  and  $X^3$  is a hydrogen atom, a halogen atom, an alkyl group, a haloalkyl group, an alkoxy group, an alkylthio group, an alkylsulfinyl group, an alkylsulfonyl group, a nitro group,  $-\text{SO}_2\text{N}(\text{R}_8)\text{R}_9$ ,  
 20  $-\text{N}(\text{R}_{10})\text{SO}_2\text{R}_{11}$ ,  $-\text{CH}_2\text{S}(\text{O})\text{qR}_{12}$  or  $-\text{OSO}_2\text{R}_{13}$ , each of  $\text{R}_8$ ,  $\text{R}_9$ ,  $\text{R}_{10}$ ,  $\text{R}_{11}$ ,  $\text{R}_{12}$  and  $\text{R}_{13}$  is an alkyl group, and  $q$  is an integer of from 0 to 2.

25



(3) The pyrazole compound or its salt according to Item 2, wherein A is  $-\text{SO}_2-$ ,  $-\text{CH}_2-$  or  $-\text{CH}_2\text{CO}-$ , each of  $\text{X}^1$ ,  $\text{X}^2$  and  $\text{X}^3$  is a hydrogen atom, a halogen atom, an alkyl group, a haloalkyl group, an alkoxy group, an alkylthio group, an alkylsulfinyl group, an alkylsulfonyl group or a nitro group.

(4) The pyrazole compound or its salt according to Item 3, wherein  $\text{X}^1$  is an alkylthio group, an alkylsulfinyl group or an alkylsulfonyl group, and each of  $\text{X}^2$  and  $\text{X}^3$  is a hydrogen atom, a halogen atom, an alkyl group, a haloalkyl group or a nitro group.

(5) A herbicide containing the pyrazole compound or its salt as defined in Item 1, 2, 3, or 4, as an active ingredient.

(6) A method for controlling noxious weeds, which comprises applying an effective amount of the pyrazole compound or its salt as defined in Item 1, 2, 3, or 4.

(7) A method for controlling noxious weeds, which comprises applying an effective amount of the pyrazole compound or its salt as defined in Item 1, 2, 3, or 4 to an upland field.

(8) A method for controlling noxious weeds, which comprises applying an effective amount of the pyrazole compound or its salt as defined in Item 1, 2, 3, or 4 to a corn field.

(9) A method for controlling noxious weeds, which comprises applying an effective amount of the pyrazole

compound or its salt as defined in Item 1, 2, 3, or 4 to a wheat field.

(10) A method for controlling noxious weeds, which comprises applying an effective amount of the pyrazole compound or its salt as defined in Item 1, 2, 3, or 4 to a paddy field.

(11) A mixed herbicidal composition comprising at least one member selected from the pyrazole compound or its salt as defined in Item 1, 2, 3, or 4 and at least one member selected from active ingredient compounds of other herbicides.


(12) The compound of the above formula (II).

(13) The compound according to Item 12, wherein 1 is 0.

(14) The compound of the above formula (IV).

(15) The compound according to Item 14, wherein 1 is 0.

(16) The compound according to Item 14, wherein the formula (IV) is represented by the formula (IV'):



(IV')

wherein  $R_1$  is an alkyl group, each of  $X^1$ ,  $X^2$  and  $X^3$  is a hydrogen atom, a halogen atom, an alkyl group, a haloalkyl group, an alkoxy group, an alkylthio group, an alkylsulfinyl group, an alkylsulfonyl group, a nitro group,  $-SO_2N(R_8)R_9$ ,  $-N(R_{10})SO_2R_{11}$ ,  $-CH_2S(O)qR_{12}$  or  $-OSO_2R_{13}$ , each of  $R_8$ ,  $R_9$ ,  $R_{10}$ ,  $R_{11}$ ,  $R_{12}$  and  $R_{13}$  is an alkyl group, and  $q$  is an integer of from 0 to 2.

(17) The pyrazole compound or its salt according to Item 16, wherein each of  $X^1$ ,  $X^2$  and  $X^3$  is a hydrogen atom, a halogen atom, an alkyl group, a haloalkyl group, an alkoxy group, an alkylthio group, an alkylsulfinyl group, an alkylsulfonyl group or a nitro group.

(18) The pyrazole compound or its salt according to Item 17, wherein  $X^1$  is an alkylthio group, an alkylsulfinyl group or an alkylsulfonyl group, and each of  $X^2$  and  $X^3$  is a hydrogen atom, a halogen atom, an alkyl group, a haloalkyl group or a nitro group.

#### BEST MODE FOR CARRYING OF THE INVENTION

Now, the present invention will be described in further detail with reference to Examples. However, it should be understood that the present invention is by no means restricted to such specific Examples. Firstly, Preparation Examples for the compounds of the present invention will be described.

#### PREPARATION EXAMPLE 1

Preparation of 3-cyclopropyl-4-(4-trifluoromethyl-2-methylsulfonylbenzoyl)-5-hydroxy-1-methylpyrazole (after-

mentioned Compound No. a-11) and 3-cyclopropyl-4-(4-trifluoromethyl-2-methylsulfonylbenzoyl)-1-methyl-5-pyrazolyl p-toluene sulfonate (after-mentioned Compound No. a-12) (First method)

5        1) 1.4 g of methylhydrazine was added at room temperature to a solution having 5.53 g of tert-butyl 3-cyclopropyl-3-oxopropionate dissolved in 30 ml of tetrahydrofuran, and the mixture was reacted for about 2 hours under reflux.

10        After completion of the reaction, tetrahydrofuran was distilled off under reduced pressure to obtain 4.14 g of crude 3-cyclopropyl-5-hydroxy-1-methylpyrazole (after-mentioned Intermediate No. 1a-1).

15        The melting point of this product was from 95 to 121°C, and the NMR spectrum data are as follows.

<sup>1</sup>H-NMR δppm [Solvent: CDCl<sub>3</sub>]

0.76-0.8 (m, 2H), 0.9-0.99 (m, 2H), 1.74-1.81 (m, 1H), 3.06 (s), 3.26 (s, 3H), 4.6 (bs)

20        2) A solution having 0.41 g of sodium carbonate dissolved in 30 ml of water, was added to a solution having 1 g of 3-cyclopropyl-5-hydroxy-1-methylpyrazole obtained in the preceding step dissolved in 30 ml of toluene, followed by stirring for 5 minutes. Then, 4-trifluoromethyl-2-methylthiobenzoyl chloride  
25        preliminarily prepared by mixing and reacting under reflux for one hour, 1.52 g of 4-trifluoromethyl-2-methylthiobenzoic acid, 5 ml of thionyl chloride and a

catalytic amount of N,N-dimethylformamide, followed by removal of excess thionyl chloride, was added thereto, and the mixture was reacted at 50°C for one hour.

After completion of the reaction, the reaction mixture was cooled and put into water, and extracted with ethyl acetate. The obtained ethyl acetate layer was washed with a saturated sodium chloride aqueous solution and then dried over anhydrous sodium sulfate, and the solvent was distilled off under reduced pressure. The obtained residue was purified by silica gel column chromatography to obtain 0.8 g of oily 3-cyclopropyl-1-methyl-5-pyrazolyl 4-trifluoromethyl-2-methylthiobenzoate (after-mentioned Intermediate No. 2a-16). The NMR spectrum data of the product are as follows.

<sup>1</sup>H-NMR δppm [Solvent: CDCl<sub>3</sub>]  
0.69-0.73 (m, 2H), 0.86-0.91 (m, 2H), 1.85-1.92 (m, 1H), 2.53 (s, 3H), 3.70 (s, 3H), 5.94 (s, 1H), 7.46 (d, 1H), 7.53 (s, 1H), 8.24 (d, 1H)

3) 0.91 g of methachloroperbenzoic acid was dividedly added at room temperature to a solution having 0.75 g of 3-cyclopropyl-1-methyl-5-pyrazolyl 4-trifluoromethyl-2-methylthiobenzoate obtained in the preceding step dissolved in 30 ml of methylene chloride, and the mixture was reacted for one hour within a range of from room temperature to 40°C.

After completion of the reaction, the reaction mixture was put into water and extracted with methylene

chloride.

The obtained methylene chloride layer was washed with dilute alkali and then with water, and thereafter dried over anhydrous sodium sulfate, and methylene chloride was distilled off. The obtained residue was purified by silica gel column chromatography to obtain 0.75 g of 3-cyclopropyl-1-methyl-5-pyrazolyl 4-trifluoromethyl-2-methylsulfonylbenzoate (after-mentioned Intermediate No. 2a-5) having a melting point of from 99 to 102°C. The NMR spectrum data of the product are as follows.

$^1\text{H-NMR}$   $\delta\text{ppm}$  [Solvent:  $\text{CDCl}_3$ ]

0.73-0.77 (m, 2H), 0.86-0.94 (m, 2H), 1.87-1.93 (m, 1H), 2.05 (s, 3H), 3.74 (s, 3H), 5.95 (s, 1H), 8.0 (d, 1H), 8.06 (d, 1H), 8.47 (s, 1H)

4) A mixture comprising 0.7 g of 3-cyclopropyl-1-methyl-5-pyrazolyl 4-trifluoromethyl-2-methylsulfonylbenzoate obtained in the preceding step, 0.3 g of dry potassium carbonate, 25 ml of toluene and 5 ml of N,N-dimethylformamide, was reacted for one hour under an azeotropic dehydration condition using a Dean-Stark azeotropic dehydration apparatus.

After completion of the reaction, the reaction mixture was cooled and put into water, and the aqueous layer was washed with ethyl acetate. The aqueous layer was acidified with concentrated hydrochloric acid and extracted with ethyl acetate. The obtained ethyl acetate layer was washed with a saturated sodium chloride aqueous

solution and dried over anhydrous sodium sulfate. Then, ethyl acetate was distilled off under reduced pressure to obtain the desired product 3-cyclopropyl-4-(4-trifluoromethyl-2-methylsulfonylbenzoyl)-5-hydroxy-1-methylpyrazole (after-mentioned Compound No. a-11) as a viscous crude product. The NMR spectrum data of this product are as follows.

$^1\text{H-NMR}$   $\delta\text{ppm}$  [Solvent:  $\text{CDCl}_3$ ]

0.42-0.45 (m, 2H), 0.72-0.81 (m, 2H), 0.95-1.05 (m, 1H),  
10 3.34 (s, 3H), 3.67 (s, 3H), 7.73 (d, 1H), 8.0 (d, 1H), 8.4 (s, 1H)

The melting point of 3-cyclopropyl-4-(4-trifluoromethyl-2-methylsulfonylbenzoyl)-5-hydroxy-1-methylpyrazole as the above-mentioned viscous crude  
15 product, was from 83 to 93°C.

5) 0.155 g of p-toluene sulfonyl chloride was added to a mixture comprising 0.3 g of 3-cyclopropyl-4-(4-trifluoromethyl-2-methylsulfonylbenzoyl)-5-hydroxy-1-methylpyrazole obtained in the preceding step, 0.118 g of  
20 dry potassium carbonate, 0.002 g of tetraethyl ammonium bromide, 20 ml of toluene and 5 ml of N,N-dimethylformamide, and the mixture was reacted for about one hour within a range of from 40 to 50°C with stirring.

After completion of the reaction, the reaction  
25 mixture was put into water and extracted with ethyl acetate. The obtained ethyl acetate layer was washed with water and further with a saturated sodium chloride

aqueous solution and then dried over anhydrous sodium sulfate, and ethyl acetate was distilled off under reduced pressure. The obtained residue was purified by silica gel column chromatography to obtain 0.3 g of 3-cyclopropyl-4-(4-trifluoromethyl-2-methylsulfonylbenzoyl)-1-methyl-5-pyrazolyl p-toluene sulfonate (after-mentioned Compound No. a-12) as a viscous desired product. The NMR spectrum data of this product are as follows.

<sup>1</sup>H-NMR δppm [Solvent: CDCl<sub>3</sub>]  
0.52-0.56 (m, 2H), 0.8-0.84 (m, 2H), 1.48-1.55 (m, 1H), 2.47 (s, 3H), 3.32 (s, 3H), 3.65 (s, 3H), 7.37 (d, 2H), 7.58 (d, 1H), 7.82 (d, 2H), 7.89 (d, 1H), 8.28 (s, 1H)

The melting point of the above viscous 3-cyclopropyl-4-(4-trifluoromethyl-2-methylsulfonylbenzoyl)-1-methyl-5-pyrazolyl p-toluene sulfonate was from 67 to 70°C.

#### PREPARATION EXAMPLE 2

Preparation of 3-cyclopropyl-4-(4-trifluoromethyl-2-methylsulfonylbenzoyl)-5-hydroxy-1-methylpyrazole (after-mentioned Compound No. a-11) and 3-cyclopropyl-4-(4-trifluoromethyl-2-methylsulfonylbenzoyl)-1-methyl-5-pyrazolyl p-toluene sulfonate (after-mentioned compound No. a-12) (Second method)

1) A mixture comprising 3.88 g of 3-cyclopropyl-1-methyl-5-pyrazolyl 4-trifluoromethyl-2-methylsulfonylbenzoate (after-mentioned Intermediate No. 2a-5), 1.52 g of dry potassium carbonate, 100 ml of



toluene and 20 ml of N,N-dimethylformamide, was reacted for one hour under an azeotropic dehydration condition using a Dean-Stark azeotropic dehydration apparatus.

After completion of the reaction, the reaction  
5 mixture was cooled and put into water, followed by liquid separation. The obtained aqueous layer was acidified with concentrated hydrochloric acid and extracted with ethyl acetate. The obtained ethyl acetate layer was washed with water and then with a saturated sodium  
10 chloride aqueous solution and dried over anhydrous sodium sulfate. Then, ethyl acetate was distilled off under reduced pressure to obtain 3.88 g of viscous 3-cyclopropyl-4-(4-trifluoromethyl-2-methylsulfonylbenzoyl)-5-hydroxy-1-methylpyrazole as a  
15 crude product. This product was left to stand to sufficiently remove the solvent to obtain crystals of 3-cyclopropyl-4-(4-trifluoromethyl-2-methylsulfonylbenzoyl)-5-hydroxy-1-methylpyrazole (after-mentioned Compound No. a-11) as the desired product,  
20 having a melting point of from 153 to 157°C.

2) 0.36 g of p-toluene sulfonyl chloride was added to a mixture comprising 0.7 g of crystals of 3-cyclopropyl-4-(4-trifluoromethyl-2-methylsulfonylbenzoyl)-5-hydroxy-1-methylpyrazole obtained in the preceding step, 0.27 g  
25 of dry potassium carbonate, 0.005 g of tetraethylammonium bromide, 20 ml of toluene and 4 ml of N,N-dimethylformamide, and the mixture was reacted for about

1.5 hours within a range of from 40 to 50°C with stirring.

After completion of the reaction, the reaction mixture was put into water and extracted with ethyl acetate. The obtained ethyl acetate layer was washed with a saturated sodium chloride aqueous solution and dried over anhydrous sodium sulfate. Then, ethyl acetate was distilled off under reduced pressure. The obtained residue was purified by silica gel column chromatography to obtain 0.7 g of crystals of 3-cyclopropyl-4-(4-trifluoromethyl-2-methylsulfonylbenzoyl)-1-methyl-5-pyrazolyl p-toluene sulfonate (after-mentioned Compound No. a-12) as the desired product, having a melting point of from 135 to 138°C.

#### 15 PREPARATION EXAMPLE 3

Preparation of 3-cyclopropyl-4-(2,4-dichloro-3-methylbenzoyl)-5-hydroxy-1-methylpyrazole (after-mentioned Compound No. a-8)

1) 2.76 g of 3-cyclopropyl-5-hydroxy-1-methylpyrazole (after-mentioned Intermediate No. 1a-1) and 3.9 g of 2,6-dichlorotoluene were charged into 30 ml of 1,2-dichloroethane, and 6.7 g of dry aluminum chloride was dividedly added thereto at a temperature of not higher than 50°C with stirring. After the addition, stirring was continued for from 10 to 15 minutes within a range of from 35 to 40°C. Then, a solution having 4.0 g of carbon tetrachloride dissolved in 4 ml of 1,2-dichloroethane,

was dropwise added thereto at the same temperature. After completion of the dropwise addition, the mixture was reacted for 1.5 hours at a temperature of from 40 to 45°C.

5       After completion of the reaction, the reaction mixture was put into 150 ml of ice water to separate a 1,2-dichloroethane layer.

2) 0.5 ml of water was added thereto, and the mixture was heated to 50°C, whereupon 3.5 ml of concentrated  
10       sulfuric acid was gradually dropwise added thereto. After completion of the dropwise addition, the mixture was reacted for 1.5 hours under reflux.

After completion of the reaction, the reaction mixture was left to cool, and 150 ml of water was added  
15       thereto, followed liquid separation. The obtained 1,2-dichloroethane layer was washed with water and then extracted with an alkaline solution having 3.5 g of sodium hydroxide dissolved in 100 ml of water. Then, 50% sulfuric acid was added thereto to make the liquid weakly  
20       acidic and extracted with methylene chloride. The obtained methylene chloride layer was dried over anhydrous sodium sulfate, and methylene chloride was distilled off under reduced pressure to obtain 3.5 g of the desired product having a melting point of from 112 to  
25       115°C. The NMR spectrum data of this product are as follows.

$^1\text{H-NMR}$   $\delta$ ppm [Solvent:  $\text{CDCl}_3$ ]

0.66-0.71 (m,2H), 0.93-0.99 (m,2H), 1.15-1.22 (m,1H),  
2.72 (s,3H), 3.89 (s,3H), 7.34 (d,1H), 7.57 (d,1H)

Preparation Example 4

Preparation of 4-(2,4-dichlorobenzoyl)-3-cyclopropyl-1-  
5 ethyl-5-hydroxypyrazole (after-mentioned Compound No. a-  
18)

1) A solution having 0.87 g of dry hydrazine  
dissolved in 5 ml of dry tetrahydrofuran, was added to a  
solution having 5 g of tert-butyl 3-cyclopropyl-3-  
10 oxopropionate dissolved in 30 ml of dry tetrahydrofuran,  
and the mixture was reacted for one hour under reflux.

After completion of the reaction, tetrahydrofuran,  
etc. were distilled off under reduced pressure to obtain  
3.3 g of 3-cyclopropyl-5-hydroxypyrazole (after-mentioned  
15 Intermediate No. 3-1) having a melting point of from 213  
to 217°C. The NMR spectrum data of this product are as  
follows.

<sup>1</sup>H-NMR δppm [Solvent: heavy MDSO]

0.57-0.61 (m,2H), 0.81-0.86 (m,2H), 1.70-1.77 (m,1H), 5.1  
20 (s,1H), 10.16 (bs,1H)

2) 1.61 g of 3-cyclopropyl-5-hydroxypyrazole obtained  
in the preceding step was mixed with a solution having  
1.89 g of dry potassium carbonate dissolved in 20 ml of  
hexamethylphosphoric triamide, and the mixture was cooled  
25 within a range of from 0 to 2°C. Then, iodoethane was  
dropwise added thereto within a range of from 0 to 5°C  
over a period of about 15 minutes. Then, the mixture was

reacted for one hour at the same temperature and then further reacted for one hour within a range of from room temperature to 40°C.

- 3) 2.72 g of 2,4-dichlorobenzoyl chloride was added thereto at room temperature, and the mixture was reacted for 0.5 hour at the same temperature and further reacted for 0.5 hour at 40°C.

After completion of the reaction, the reaction mixture was put into water and extracted with toluene. The obtained toluene layer was thoroughly washed with water and then with a saturated sodium chloride aqueous solution and then dried over anhydrous sodium sulfate. Then, toluene was distilled off under reduced pressure, and the obtained residue was purified by silica gel column chromatography to obtain 1.2 g of 3-cyclopropyl-1-ethyl-5-pyrazolyl 2,4-dichlorobenzoate (after-mentioned Intermediate No. 2a-7) having a melting point of from 61 to 63°C. The NMR spectrum data of this product are as follows.

<sup>1</sup>H-NMR δppm [Solvent: CDCl<sub>3</sub>]  
0.69-0.73 (m, 2H), 0.87-0.9 (m, 2H), 1.4 (t, 3H), 1.85-1.92 (m, 1H), 4.02-4.08 (q, 2H), 5.92 (s, 1H), 7.39 (d, 1H), 7.55 (s, 1H), 7.94 (d, 1H)

- 4) Using 1.1 g of 3-cyclopropyl-1-ethyl-5-pyrazolyl 2,4-dichlorobenzoate obtained in the preceding step, 0.843 g of the desired product having a melting point of from 74 to 77°C was obtained in the same manner as Step

4) in Preparation Example 1.

PREPARATION EXAMPLE 5

Preparation of 3-cyclopropyl-4-(4-trifluoromethyl-2-methylsulfonylbenzoyl)-1-methyl-5-pyrazolyl benzene

5 sulfonate (after-mentioned Compound No. a-27)

1) 0.57 g of 3-cyclopropyl-1-methyl-5-pyrazolyl 4-trifluoromethyl-2-methylsulfonyl benzoate (after-mentioned Intermediate No. 2a-5), 20 ml of toluene and 1 ml of N,N-dimethylformamide were charged into an  
10 Erlenmeyer flask, and 0.11 g of potassium carbonate was added thereto. The mixture was reacted for 15 hours under an azeotropic dehydration condition to obtain a reaction mixture containing a potassium salt of 3-cyclopropyl-4-(4-trifluoromethyl-2-  
15 methylsulfonylbenzoyl)-5-hydroxy-1-methylpyrazole.

2) The reaction mixture obtained in the preceding step was left to cool, and 0.1 g of tetraethylammonium chloride and 0.1 g of potassium iodide were added thereto. Then, 0.27 g of benzene sulfonyl chloride was  
20 added thereto. The mixture was reacted for 5.5 hours at 55°C with stirring.

After completion of the reaction, the reaction mixture was put into water and extracted with ethyl acetate. Then, the extract was washed with water. The  
25 obtained organic layer was dried over anhydrous sodium sulfate, then concentrated and thereafter purified by silica gel column chromatography to obtain 0.49 g of the

desired product having a melting point of from 175 to 178°C. The NMR spectrum data of this product are as follows.

<sup>1</sup>H-NMR δppm [Solvent: CDCl<sub>3</sub>]

5 0.46-0.05 (m, 2H), 0.73-0.81 (m, 2H), 1.33-1.41 (m, 1H),  
3.27 (s, 3H), 3.63 (s, 3H), 7.53-7.58 (m, 3H), 7.7 (t, 1H),  
7.85 (d, 1H), 7.96 (d, 2H), 8.27 (s, 1H)

#### PREPARATION EXAMPLE 6

Preparation of 3-cyclopropyl-4-(4-trifluoromethyl-2-  
10 methysulfonylbenzoyl)-1-methyl-5-pyrazolyl n-  
propanesulfonate (after-mentioned Compound No. a-89)

A mixture comprising 0.4 g of 3-cyclopropyl-4-(4-trifluoromethyl-2-methysulfonylbenzoyl)-5-hydroxy-1-methylpyrazole (after-mentioned Compound No. a-11), 20 ml  
15 of toluene, 5 ml of N,N-dimethylformamide, 5 mg of tetraethylammonium bromide and 0.16 g of n-propanesulfonyl chloride, was reacted for about 12 hours at 40°C with stirring.

After completion of the reaction, the reaction  
20 mixture was put into water and extracted with ethyl acetate. The obtained ethyl acetate layer was washed with a saturated sodium chloride aqueous solution and then dried over anhydrous sodium sulfate. Then, ethyl acetate was distilled off under reduced pressure. The  
25 obtained residue was purified by silica gel column chromatography to obtain 0.34 g of the desired product having a melting point of from 128 to 131°C. The NMR

spectrum data of this product are as follows.

$^1\text{H}$ -NMR  $\delta$ ppm [Solvent:  $\text{CDCl}_3$ ]

0.43-0.51 (m, 2H), 0.78-0.82 (m, 2H), 1.12 (t, 3H), 1.1-1.2  
(m, 1H), 2.0-2.1 (m, 2H), 3.33 (s, 3H), 3.53 (t, 2H), 3.82  
5 (s, 3H), 7.70 (d, 1H), 7.96 (d, 1H), 8.38 (s, 1H)

#### PREPARATION EXAMPLE 7

Preparation of 5-benzyloxy-3-cyclopropyl-4-(4-  
trifluoromethyl-2-methylsulfonylbenzoyl)-1-methylpyrazole  
(after-mentioned Compound No. a-94)

10 0.14 g of benzyl chloride was added to a mixture  
comprising 0.4 g of 3-cyclopropyl-4-(4-trifluoromethyl-2-  
methylsulfonylbenzoyl)-5-hydroxy-1-methylpyrazole (after-  
mentioned Compound No. a-11), 0.16 g of dry potassium  
carbonate, 5 mg of benzyltriethylammonium chloride, 5 mg  
15 of potassium iodide, 20 ml of toluene and 5 ml of N,N-  
dimethylformamide, and the mixture was reacted for 24  
hours within a range of from 50 to 70°C with stirring.

After completion of the reaction, the reaction  
mixture was put into water and extracted with ethyl  
20 acetate. The obtained ethyl acetate layer was washed  
with a saturated sodium chloride aqueous solution and  
then dried over anhydrous sodium sulfate. Then, ethyl  
acetate was distilled off under reduced pressure. The  
obtained residue was purified by silica gel column  
25 chromatography to obtain 0.25 g of the desired product  
having a melting point of from 154 to 157°C. The NMR  
spectrum data of this product are as follows.



<sup>1</sup>H-NMR δppm [Solvent: CDCl<sub>3</sub>]

0.68-0.71 (m, 2H), 0.85-0.88 (m, 2H), 1.8-2.0 (m, 1H), 3.35 (s, 3H), 3.42 (s, 3H), 5.00 (s, 2H), 7.11-7.12 (m, 2H), 7.26-7.30 (m, 3H), 7.58-7.60 (d, 1H), 7.58-7.88 (d, 1H), 8.34 (s, 1H)

PREPARATION EXAMPLE 8

Preparation of 5-(2-chloro-2-propenyloxy)-3-cyclopropyl-4-(4-trifluoromethyl-2-methylsulfonylbenzoyl)-1-methylpyrazole (after-mentioned Compound No. a-213)

10 A mixture comprising 0.776 g of 3-cyclopropyl-4-(4-trifluoromethyl-2-methylsulfonylbenzoyl)-5-hydroxy-1-methylpyrazole (after-mentioned Compound No. a-11), 30 ml of toluene, 4 ml of N,N-dimethylformamide, 5 mg of tetraethylammonium bromide and 0.245 g of 2,3-  
15 dichloropropene, was reacted for 1 hour at room temperature, then reacted for 4 hours at a temperature of from 60 to 80°C with stirring.

After completion of the reaction, the reaction mixture was put into water and extracted with ethyl  
20 acetate. The obtained ethyl acetate layer was washed with a saturated sodium chloride aqueous solution and then dried over anhydrous sodium sulfate. Then, ethyl acetate was distilled off. The obtained residue was purified by silica gel column chromatography to obtain  
25 0.65 g of the desired product having a melting point of from 180 to 111°C. The NMR spectrum data of this product are as follows.

<sup>1</sup>H-NMR δppm [Solvent: CDCl<sub>3</sub>]

0.59-0.61 (m, 2H), 0.84-0.86 (m, 2H), 1.6-1.7 (m, 1H), 3.38 (s, 3H), 3.67 (s, 3H), 4.68 (s, 2H), 5.37-5.4 (d, 2H), 7.62 (d, 1H), 7.93 (d, 1H), 8.38 (s, 1H)

5 PREPARATION EXAMPLE 9

Preparation of 3-cyclopropyl-1-methyl-4-(2-methylthio-4-trifluoromethylbenzoyl)-5-hydroxypyrazole (after-mentioned Compound No. a-82), 3-cyclopropyl-1-methyl-4-(2-methylthio-4-trifluoromethylbenzoyl)-5-pyrazolyl p-toluene sulfonate (after-mentioned Compound No. a-72) and 3-cyclopropyl-4-(4-trifluoromethyl-2-methylsulfonylbenzoyl)-1-methyl-5-pyrazolyl p-toluene sulfonate (after-mentioned Compound No. a-12)

10

1) Into a 200 ml autoclave, 1.59 g of 4-iodo-3-methylthiobenzotrifluoride prepared in accordance with the following Preparation Example 10, 1.38 g of 3-cyclopropyl-5-hydroxy-1-methylpyrazole (after-mentioned Intermediate No. 1a-1), 0.5 g of triethylamine, 3.1 g of potassium carbonate, 0.22 g of palladium (II) bis(triphenylphosphine) dichloride and 40 ml of dioxane were put and sealed, and the interior of the autoclave was flushed with carbon monoxide (pressure: 65 kg/cm<sup>2</sup>), followed by a reaction at 140°C for 8 hours. After completion of the reaction, the solvent was distilled off, and the residue was dissolved in water, and then insoluble matters were filtered off. The filtrate was washed with dichloromethane. The washed product was

15

20

25

acidified (pH=1) with concentrated hydrochloric acid and extracted with dichloromethane. The obtained extract solution was dried over anhydrous sodium sulfate, and the solvent was distilled off to obtain 1.59 g of 3-cyclopropyl-1-methyl-4-(2-methylthio-4-trifluoromethylbenzoyl)-5-hydroxypyrazole (after-mentioned Compound No. a-82) as a reddish brown solid.

2) 1.59 g of 3-cyclopropyl-1-methyl-4-(2-methylthio-4-trifluoromethylbenzoyl)-5-hydroxypyrazole obtained in the preceding step was mixed, without purification, with 20 ml of toluene, 4 ml of N,N-dimethylformamide, 0.94 g of p-toluene sulfonyl chloride and 0.34 g of potassium carbonate, and the mixture was reacted at 60°C for 3 hours. After completion of the reaction, water was added to the reaction mixture, and the mixture was extracted with ethyl acetate. The extract solution was dried over anhydrous sodium sulfate, and the solvent was distilled off. The obtained residue was purified by silica gel column chromatography (developing solvent: ethyl acetate/hexane=1/4) to obtain 0.53 g of 3-cyclopropyl-1-methyl-4-(2-methylthio-4-trifluoromethylbenzoyl)-5-pyrazolyl p-toluene sulfonate (after-mentioned Compound No. a-72). The NMR spectrum data of this product are as follows.

<sup>1</sup>H-NMR δppm [Solvent: CDCl<sub>3</sub>]  
0.79 (m,2H), 0.90 (m,2H), 1.97 (m,1H), 2.39 (s,3H), 2.47 (s,3H), 7.23 (d,2H), 7.32 (d,1H), 7.48 (s,1H), 7.49

(d,1H), 7.53 (d,2H)

3). 0.46 g of 3-cyclopropyl-1-methyl-4-(2-methylthio-4-trifluoromethylbenzoyl)-5-pyrazolyl p-toluene sulfonate obtained in the preceding step was dissolved in 10 ml of dichloromethane, and 0.47 g of 85% methachloroperbenzoic acid was added thereto under cooling with ice. Then, the mixture was returned to room temperature and reacted over night with stirring. After completion of the reaction, an aqueous sodium hydrogen carbonate solution was added to the reaction mixture, and the mixture was extracted with dichloromethane. The extract layer was dried over anhydrous sodium sulfate, and the solvent was distilled off. The obtained residue was purified by silica gel column chromatography (developing solvent: ethyl acetate/hexane=3/7) to obtain 0.49 g of 3-cyclopropyl-4-(4-trifluoromethyl-2-methylsulfonylbenzoyl)-1-methyl-5-pyrazolyl p-toluene sulfonate (after-mentioned Compound No. a-12).

#### PREPARATION EXAMPLE 10

##### 20 Preparation of 4-iodo-3-methylthiobenzotrifluoride

1) 123.85 g of sodium iodide was added to a solution having 42.23 g of 4-chloro-3-nitrobenzotrifluoride dissolved in 200 ml of N,N-dimethylformamide, and the mixture was reacted at 140°C for 17 hours.

25 After completion of the reaction, the reaction mixture was put into water and extracted with ethyl ether. The ethyl ether layer was washed with water and

then dried over anhydrous sodium sulfate. Then, ethyl ether was distilled off. The obtained residue was purified by silica gel column chromatography to obtain 44.15 g of 4-iodo-3-nitrobenzotrifluoride. The NMR spectrum data of this product are as follows.

$^1\text{H-NMR}$   $\delta\text{ppm}$  [Solvent:  $\text{CDCl}_3$ ]

7.52 (dd,1H), 8.11 (s,1H), 8.22 (d,1H)

2) A solution having 30 g of 4-iodo-3-nitrobenzotrifluoride obtained in the preceding step dissolved in 300 ml of acetic acid, was heated, and 26.43 g of reduced iron was added thereto over a period of 15 minutes at a temperature of from 85 to 95°C. Then, the mixture was reacted for further 5 minutes at the same temperature.

After completion of the reaction, the reaction mixture was cooled with ice, and insoluble matters were filtered off using celite. The filtration cake was thoroughly washed with ethyl acetate, and the washing liquid and the filtrate were mixed, followed by washing with water for 5 times. The obtained ethyl acetate layer was dried over anhydrous sodium sulfate, and ethyl acetate was distilled off under reduced pressure. The obtained residue was purified by silica gel column chromatography to obtain 25.52 g of oily 3-amino-4-iodobenzotrifluoride. The NMR spectrum data of this product are as follows.

$^1\text{H-NMR}$   $\delta\text{ppm}$  [Solvent:  $\text{CDCl}_3$ ]

6.70 (dd,1H), 6.93 (d,1H), 7.73 (d,1H)

3) To a solution containing a part (5.1 g) of 3-amino-4-iodobenzotrifluoride obtained in the preceding step, 16.75 g of dimethyldisulfide and 80 ml of  
5 chloroform, a solution having the rest (20.42 g) of 3-amino-4-iodobenzotrifluoride obtained in the preceding step dissolved in 20 ml of chloroform and 11.92 g of tert-butyl nitrite, were simultaneously dropwise added at a temperature of from 25 to 30°C. After completion of  
10 the dropwise addition, the mixture was reacted at room temperature for 16 hours.

After completion of the reaction, 200 ml of methylene chloride was added to the reaction mixture, and the mixture was washed with an aqueous hydrochloric acid  
15 solution with pH 1 to 2. Then, the methylene chloride layer was washed with water and dried over anhydrous sodium sulfate. Then, methylene chloride and chloroform were distilled off under reduced pressure. The obtained residue was purified by silica gel column chromatography  
20 to obtain 19.89 g of the desired product as an oily substance. The NMR spectrum data of this product are as follows.

$^1\text{H-NMR}$   $\delta$ ppm [Solvent:  $\text{CDCl}_3$ ]

2.51 (s,3H), 7.08 (dd,1H), 7.26 (d,1H), 7.90 (dd,1H)

25 Other compounds of the present invention can be prepared in accordance with the above described Preparation Examples or the above described various

processes for producing the compounds of the present invention. Typical examples of the intermediate compound represented by the above formula (II) will be shown in Table 1, typical examples of the intermediate compound represented by the above formula (IV) will be presented in Tables 2a and 2b, typical examples of the intermediate compound represented by the above formula (IX) will be presented in Table 3, and typical examples of the compound of the present invention represented by the above formula (I) will be presented in Tables 4a and 4b.

62















Table 1

Inter- mediate No.	R <sub>1</sub>	(Z) 	Physical properties
1 a - 1	CH <sub>3</sub>		m.p. 95 - 121 °C
1 a - 2	CH <sub>2</sub> CH <sub>3</sub>		
1 a - 3	n-C <sub>3</sub> H <sub>7</sub>		
1 a - 4	n-C <sub>4</sub> H <sub>9</sub>		
1 a - 5	CH(CH <sub>3</sub> ) <sub>2</sub>		
1 b - 1	CH <sub>3</sub>	CH <sub>3</sub> 	
1 b - 2	CH <sub>2</sub> CH <sub>3</sub>	CH <sub>3</sub> 	
1 b - 3	n-C <sub>3</sub> H <sub>7</sub>	CH <sub>3</sub> 	
1 b - 4	n-C <sub>4</sub> H <sub>9</sub>	CH <sub>3</sub> 	
1 b - 5	CH(CH <sub>3</sub> ) <sub>2</sub>	CH <sub>3</sub> 	



63




Table 2a

Inter- mediate No.	R <sub>1</sub>	(X) <sub>n</sub>	Physical properties
2 a - 1	CH <sub>3</sub>		
2 a - 2	CH <sub>3</sub>		m.p. 84-87°C
2 a - 3	CH <sub>3</sub>		m.p. 148-150°C
2 a - 4	CH <sub>3</sub>		
2 a - 5	CH <sub>3</sub>		m.p. 99-102°C
2 a - 6	CH <sub>3</sub>		
2 a - 7	CH <sub>2</sub> CH <sub>3</sub>		m.p. 61-63°C
2 a - 8	CH <sub>3</sub>		

Table 2a (continued)











Inter- mediate No.	R <sub>1</sub>	(X) <sub>n</sub> 	Physical properties
2 a - 9	CH <sub>2</sub> CH <sub>3</sub>		m.p. 96-99°C
2 a - 10	CH <sub>3</sub>		Viscous
2 a - 11	CH <sub>3</sub>		
2 a - 12	CH <sub>2</sub> CH <sub>3</sub>		
2 a - 13	n-C <sub>3</sub> H <sub>7</sub>		
2 a - 14	CH <sub>2</sub> CH <sub>3</sub>		m.p. 94-97°C
2 a - 15	CH(CH <sub>3</sub> ) <sub>2</sub>		
2 a - 16	CH <sub>3</sub>		Oily
2 a - 17	CH <sub>3</sub>		m.p. 136-140 °C

Table 2a (continued)











Inter- mediate	R <sub>1</sub>	(X) <sub>n</sub> 	Physical properties
2 a - 18	CH <sub>3</sub>		
2 a - 19	CH <sub>3</sub>		m.p. 90—93°C
2 a - 20	CH <sub>2</sub> CH <sub>3</sub>		
2 a - 21	CH <sub>3</sub>		Viscous
2 a - 22	CH <sub>3</sub>		Viscous
2 a - 23	CH <sub>3</sub>		Viscous
2 a - 24	CH <sub>3</sub>		m.p. 146 — 149 °C
2 a - 25	CH <sub>3</sub>		m.p. 125 — 130 °C
2 a - 26	CH <sub>3</sub>		

Table 2a (continued)










Inter- mediate No.	R <sub>1</sub>	(X) <sub>n</sub> 	Physical properties
2 a - 27	CH <sub>3</sub>		
2 a - 28	CH <sub>3</sub>		m.p. 134 - 136 °C
2 a - 29	CH <sub>3</sub>		
2 a - 30	CH <sub>3</sub>		
2 a - 31	CH <sub>3</sub>		
2 a - 32	CH <sub>3</sub>		
2 a - 33	CH <sub>3</sub>		
2 a - 34	CH <sub>3</sub>		

Table 2a (continued)











Inter- mediate No.	R <sub>1</sub>		Physical properties
2 a - 35	CH <sub>3</sub>		
2 a - 36	CH <sub>3</sub>		
2 a - 37	CH <sub>3</sub>		
2 a - 38	CH <sub>3</sub>		
2 a - 39	CH <sub>3</sub>		
2 a - 40	CH <sub>3</sub>		
2 a - 41	CH <sub>3</sub>		
2 a - 42	CH <sub>3</sub>		
2 a - 43	CH <sub>3</sub>		m.p. 118 - 122 °C

Table 2a (continued)










Inter- mediate No.	R <sub>1</sub>	(X) <sub>n</sub> 	Physical properties
2 a - 44	CH <sub>3</sub>		
2 a - 45	CH <sub>3</sub>		Viscous
2 a - 46	CH <sub>3</sub>		
2 a - 47	CH <sub>3</sub>		
2 a - 48	CH <sub>3</sub>		
2 a - 49	CH <sub>3</sub>		
2 a - 50	CH <sub>3</sub>		
2 a - 51	CH <sub>3</sub>		

Table 2a (continued)

















Inter- mediate No.	R <sub>1</sub>		Physical properties
2 a - 52	CH <sub>3</sub>		
2 a - 53	CH <sub>3</sub>		
2 a - 54	CH <sub>3</sub>		
2 a - 55	CH <sub>3</sub>		
2 a - 56	CH <sub>3</sub>		
2 a - 57	CH <sub>3</sub>		
2 a - 58	CH <sub>3</sub>		
2 a - 59	CH <sub>3</sub>		
2 a - 60	CH <sub>3</sub>		

Table 2a (continued)

Inter- mediate No.	R <sub>1</sub>	(X) <sub>n</sub> 	Physical properties
2 a - 61	CH <sub>3</sub>	SC <sub>2</sub> H <sub>5</sub> (iso) 	
2 a - 62	CH <sub>3</sub>	SO <sub>2</sub> C <sub>2</sub> H <sub>5</sub> (iso) 	
2 a - 63	CH <sub>3</sub>	SO <sub>2</sub> C <sub>2</sub> H <sub>5</sub> (iso) 	
2 a - 64	CH <sub>3</sub>		
2 a - 65	CH <sub>3</sub>		





















Table 2b

Inter- mediate No.	R <sub>1</sub>	(Z) 		Physical properties
2 b - 1	CH <sub>3</sub>	CH <sub>3</sub> 	SO <sub>2</sub> CH <sub>3</sub> 	Viscous
2 b - 2	CH <sub>2</sub> CH <sub>3</sub>	CH <sub>3</sub> 	SO <sub>2</sub> CH <sub>3</sub> 	
2 b - 3	CH <sub>3</sub>	CH <sub>3</sub> 	SO <sub>2</sub> CH <sub>3</sub> 	
2 b - 4	CH <sub>3</sub>	CH <sub>3</sub> 	CF <sub>3</sub> 	
2 b - 5	CH <sub>3</sub>	CH <sub>3</sub> 	SCH <sub>3</sub> 	
2 b - 6	CH <sub>3</sub>	CH <sub>2</sub> 	Cl CH <sub>3</sub> 	



... (IX)

Table 3

Inter- mediate No.	(Z)	Physical properties
3 - 1		m.p. 213 - 217 °C
3 - 2	CH <sub>3</sub>	

73




Table 4a

Compound No.	R <sub>1</sub>	R <sub>2</sub>	(X) <sub>n</sub>	Physical properties
a - 1	CH <sub>3</sub>	H		m.p. 131-133°C
a - 2	CH <sub>3</sub>	-SO <sub>2</sub> -		Refractive index 13.4 n <sub>D</sub> 1.5779
a - 3	CH <sub>3</sub>	H		m.p. 65-70°C
a - 4	CH <sub>3</sub>	-SO <sub>2</sub> -		m.p. 130-133°C
a - 5	CH <sub>3</sub>	H		m.p. 163-166°C
a - 6	CH <sub>3</sub>	-SO <sub>2</sub> -		m.p. 172-174°C
a - 7	CH <sub>3</sub>	-CH <sub>2</sub> C(=O)-		m.p. 145-147°C
a - 8	CH <sub>3</sub>	H		m.p. 112-115°C

74  
Table 4a (continued)

















Compound No.	R <sub>1</sub>	R <sub>2</sub>	(X) <sub>n</sub>	Physical properties
a - 9	CH <sub>3</sub>	-SO <sub>2</sub> -  -CH <sub>3</sub>		m.p. 115 - 118°C
a - 10	CH <sub>3</sub>	-CH <sub>2</sub> -C(=O)-  -CH <sub>3</sub>		m.p. 126 - 129°C
a - 11	CH <sub>3</sub>	H		m.p. 153 - 157°C
a - 12	CH <sub>3</sub>	-SO <sub>2</sub> -  -CH <sub>3</sub>		m.p. 135 - 138°C
a - 13	CH <sub>3</sub>	-CH <sub>2</sub> -C(=O)- 		m.p. 124 - 127°C
a - 14	CH <sub>3</sub>	-C(=O)CH <sub>3</sub>		m.p. 112 - 115°C
a - 15	CH <sub>3</sub>	H		m.p. 115 - 122°C
a - 16	CH <sub>3</sub>	-SO <sub>2</sub> -  -CH <sub>3</sub>		m.p. 146 - 148°C
a - 17	CH <sub>3</sub>	-CH <sub>2</sub> -C(=O)- 		Viscous
a - 18	CH <sub>2</sub> CH <sub>3</sub>	H		m.p. 74 - 77°C

Table 4a (continued)

Compound No.	R <sub>1</sub>	R <sub>2</sub>	(X) <sub>n</sub>	Physical properties
a - 19	CH <sub>2</sub> CH <sub>3</sub>			Viscous
a - 20	CH <sub>3</sub>			m.p. 181 - 183°C
a - 21	CH <sub>2</sub> CH <sub>3</sub>	H		m.p. 158 - 161°C
a - 22	CH <sub>2</sub> CH <sub>3</sub>			m.p. 116 - 118°C
a - 23	CH <sub>2</sub> CH <sub>3</sub>			m.p. 146 - 148°C
a - 24	CH <sub>3</sub>	H		m.p. 111 - 114°C
a - 25	CH <sub>3</sub>			Refractive index n <sub>D</sub> <sup>20</sup> 1.6001
a - 26	CH <sub>3</sub>	H		m.p. 140 - 145°C
a - 27	CH <sub>3</sub>			m.p. 175 - 178°C

Table 4a (continued)




















Compound No.	R <sub>1</sub>	R <sub>2</sub>		Physical properties
a - 28	CH <sub>3</sub>			
a - 29	CH <sub>3</sub>			
a - 30	CH <sub>3</sub>			
a - 31	CH <sub>2</sub> CH <sub>3</sub>			
a - 32	CH <sub>2</sub> CH <sub>3</sub>			
a - 33	CH <sub>3</sub>			
a - 34	CH <sub>3</sub>			
a - 35	CH <sub>3</sub>			
a - 36	CH <sub>3</sub>			

Table 4a (continued)




















Compound No.	R <sub>1</sub>	R <sub>2</sub>		Physical properties
a - 37	CH <sub>3</sub>			
a - 38	CH <sub>3</sub>			
a - 39	CH <sub>3</sub>	-SO <sub>2</sub> CH <sub>3</sub>		m. p. 143 - 146°C
a - 40	CH <sub>3</sub>	-SO <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>		m. p. 127 - 130.5°C
a - 41	CH <sub>3</sub>			m. p. 117 - 120°C
a - 42	CH <sub>3</sub>			m. p. 196 - 199°C
a - 43	CH <sub>3</sub>			m. p. 157 - 160°C
a - 44	CH <sub>3</sub>			m. p. 168 - 171°C
a - 45	CH <sub>3</sub>			m. p. 157 - 160°C
a - 46	CH <sub>3</sub>			

Table 4a (continued)




















Compound No.	R <sub>1</sub>	R <sub>2</sub>	(X) <sub>n</sub>	Physical properties
a - 47	CH <sub>3</sub>	-SO <sub>2</sub> -  -CF <sub>3</sub>		m.p. 185 - 188°C
a - 48	CH <sub>3</sub>	-SO <sub>2</sub> -  -CF <sub>3</sub>		m.p. 124 - 127°C
a - 49	CH <sub>3</sub>	-SO <sub>2</sub> -  -F		m.p. 152 - 155°C
a - 50	CH <sub>3</sub>	-SO <sub>2</sub> -  -F		
a - 51	CH <sub>3</sub>	-SO <sub>2</sub> -  -CH <sub>3</sub>		m.p. 166 - 169°C
a - 52	CH <sub>3</sub>	-SO <sub>2</sub> -  -CH <sub>3</sub>		m.p. 145 - 149°C
a - 53	CH <sub>3</sub>	-SO <sub>2</sub> C <sub>4</sub> H <sub>9</sub> (n)		m.p. 142 - 145°C
a - 54	CH <sub>2</sub> CH <sub>3</sub>	-SO <sub>2</sub> - 		
a - 55	CH <sub>2</sub> CH <sub>3</sub>	-SO <sub>2</sub> -  -CH <sub>3</sub>		m.p. 156 - 159°C
a - 56	CH <sub>2</sub> CH <sub>3</sub>	-SO <sub>2</sub> -  -Cl		



Table 4a (continued)





















Compound No.	R <sub>1</sub>	R <sub>2</sub>		Physical properties
a - 57	CH <sub>2</sub> CH <sub>3</sub>			
a - 58	CH <sub>2</sub> CH <sub>3</sub>			m.p. 136 - 138°C
a - 59	CH <sub>3</sub>			viscous
a - 60	CH <sub>3</sub>			Viscous
a - 61	CH(CH <sub>3</sub> ) <sub>2</sub>			
a - 62	CH(CH <sub>3</sub> ) <sub>2</sub>			
a - 63	n-C <sub>3</sub> H <sub>7</sub>			
a - 64	CH <sub>3</sub>	-SO <sub>2</sub> CH <sub>3</sub>		
a - 65	CH <sub>3</sub>			
a - 66	CH <sub>3</sub>			

Table 4a (continued)

Compound No.	R <sub>1</sub>	R <sub>2</sub>	(X) <sub>n</sub> 	Physical properties
a - 67	CH <sub>3</sub>	-SO <sub>2</sub> -		
a - 68	CH <sub>3</sub>	-SO <sub>2</sub> -		
a - 69	CH <sub>3</sub>	-SO <sub>2</sub> -		m. p. 208 - 211°C
a - 70	CH <sub>3</sub>	-SO <sub>2</sub> -		
a - 71	n-C <sub>3</sub> H <sub>7</sub>	-SO <sub>2</sub> -		
a - 72	CH <sub>3</sub>	-SO <sub>2</sub> -		m. p. 107 - 109°C
a - 73	CH <sub>3</sub>	-SO <sub>2</sub> -		m. p. 158 - 164°C
a - 74	CH <sub>3</sub>	-SO <sub>2</sub> -		
a - 75	CH <sub>3</sub>	-SO <sub>2</sub> -		
a - 76	CH <sub>3</sub>	-SO <sub>2</sub> -		Viscous

Table 4a (continued)

















Compound No.	R <sub>1</sub>	R <sub>2</sub>		Physical properties
a - 77	CH <sub>3</sub>	H		m.p. 190 - 204°C
a - 78	CH <sub>3</sub>	-SO <sub>2</sub> CH <sub>3</sub>		m.p. 134 - 138°C
a - 79	CH <sub>3</sub>			m.p. 137 - 139°C
a - 80	CH <sub>3</sub>	H		m.p. 177 - 180°C
a - 81	CH <sub>3</sub>	H		m.p. 141 - 143°C
a - 82	CH <sub>3</sub>	H		m.p. 107 - 110°C
a - 83	CH <sub>3</sub>			Viscous
a - 84	CH <sub>3</sub>			m.p. 189 - 193°C
a - 85	CH <sub>3</sub>			m.p. 113 - 115°C
a - 86	CH <sub>3</sub>			m.p. 146 - 148°C

Table 4a (continued)

Compound No.	R <sub>1</sub>	R <sub>2</sub>	(X) <sub>n</sub>	Physical properties
a - 87	CH <sub>3</sub>	$-\text{CH}_2\text{C}(=\text{O})-\text{C}_6\text{H}_5$	$\text{SOCH}_3$ $-\text{C}_6\text{H}_4-\text{CF}_3$	m.p. 148 - 151°C
a - 88	CH <sub>3</sub>	$-\text{C}(=\text{O})-\text{C}_6\text{H}_5$	$\text{SO}_2\text{CH}_3$ $-\text{C}_6\text{H}_4-\text{CF}_3$	m.p. 137 - 141°C
a - 89	CH <sub>3</sub>	$-\text{SO}_2\text{C}_2\text{H}_5 (n)$	$\text{SO}_2\text{CH}_3$ $-\text{C}_6\text{H}_4-\text{CF}_3$	m.p. 128 - 131°C
a - 90	CH <sub>3</sub>	$-\text{SO}_2-\text{C}_6\text{H}_5$	$\text{Cl}$ $\text{CH}_3$ $-\text{C}_6\text{H}_3-\text{SO}_2\text{CH}_3$	Viscous
a - 91	CH <sub>3</sub>	$-\text{SO}_2-\text{C}_6\text{H}_4-\text{CH}_3$	$\text{Cl}$ $\text{CH}_3$ $-\text{C}_6\text{H}_3-\text{SO}_2\text{CH}_3$	m.p. 188 - 192°C
a - 92	CH <sub>3</sub>	$-\text{SO}_2-\text{C}_6\text{H}_4-\text{CH}_3$	$\text{CF}_3$ $-\text{C}_6\text{H}_4-\text{SO}_2\text{CH}_3$	Viscous
a - 93	CH <sub>3</sub>	$-\text{SO}_2-\text{C}_6\text{H}_4-\text{CH}_3$	$\text{SOCH}_3$ $-\text{C}_6\text{H}_4-\text{CF}_3$	m.p. 128 - 131°C
a - 94	CH <sub>3</sub>	$-\text{CH}_2-\text{C}_6\text{H}_5$	$\text{SO}_2\text{CH}_3$ $-\text{C}_6\text{H}_4-\text{CF}_3$	m.p. 154 - 157°C
a - 95	CH <sub>3</sub>	$-\text{CH}_2\text{CN}$	$\text{SO}_2\text{CH}_3$ $-\text{C}_6\text{H}_4-\text{CF}_3$	m.p. 135 - 140°C
a - 96	CH <sub>3</sub>	$-\text{CH}_2\text{CH}=\text{CH}_2$	$\text{SO}_2\text{CH}_3$ $-\text{C}_6\text{H}_4-\text{CF}_3$	Refractive index at 25°C n <sub>D</sub> 1.5133

Table 4a (continued)


















Compound No.	R <sub>1</sub>	R <sub>2</sub>		Physical properties
a - 97	CH <sub>3</sub>	-CH <sub>2</sub> -  -CH <sub>3</sub>		m.p. 161 - 163°C
a - 98	CH <sub>3</sub>	-CH <sub>2</sub> - 		m.p. 163 - 166°C
a - 99	CH <sub>3</sub>	-CH <sub>2</sub> C≡CH		m.p. 123 - 127°C
a - 100	CH <sub>3</sub>	-CH <sub>2</sub> CH <sub>3</sub>		m.p. 125 - 128°C
a - 101	CH <sub>3</sub>	-CH <sub>2</sub> -  -CF <sub>3</sub>		m.p. 159 - 161°C
a - 102	CH <sub>3</sub>	-SO <sub>2</sub> -  -CH <sub>3</sub>		m.p. 63 - 70°C
a - 103	CH <sub>3</sub>	-SO <sub>2</sub> C <sub>3</sub> H <sub>7</sub> (n)		m.p. 130 - 133°C
a - 104	CH <sub>3</sub>	-CH <sub>2</sub> -  -Cl		m.p. 151 - 154°C
a - 105	CH <sub>3</sub>	-CH <sub>2</sub> -  -Br		m.p. 160 - 163°C
a - 106	CH <sub>3</sub>	H		m.p. 140 - 143°C

Table 4a (continued)















Compound No.	R <sub>1</sub>	R <sub>2</sub>		Physical properties
a -107	CH <sub>2</sub> CH <sub>3</sub>	H		m.p. 109 - 114°C
a -108	CH <sub>3</sub>	-CH <sub>2</sub> - 		
a -109	CH <sub>3</sub>	-CH <sub>2</sub> - 		
a -110	CH <sub>3</sub>	-CH <sub>2</sub> - 		
a -111	CH <sub>3</sub>	-CH <sub>2</sub> OCH <sub>3</sub>		
a -112	CH <sub>3</sub>	-CH <sub>2</sub> OCH <sub>2</sub> CH <sub>3</sub>		
a -113	CH <sub>3</sub>	$\begin{array}{c} \text{O} \\ \parallel \\ -\text{CH}_2\text{COCH}_3 \end{array}$		Viscous
a -114	CH <sub>3</sub>	$\begin{array}{c} \text{O} \\ \parallel \\ -\text{CH}_2\text{COCH}_2\text{CH}_3 \end{array}$		
a -115	CH <sub>3</sub>	-CH <sub>3</sub>		
a -116	CH <sub>3</sub>	-C <sub>3</sub> H <sub>7</sub> (n)		Viscous

Table 4a (continued)












Compound No.	R <sub>1</sub>	R <sub>2</sub>		Physical properties
a -117	CH <sub>3</sub>	-C <sub>4</sub> H <sub>9</sub> (n)		
a -118	CH <sub>3</sub>	$\begin{array}{c} \text{O} \\ \parallel \\ -\text{CCH}_3 \end{array}$		m.p. 156 - 158°C
a -119	CH <sub>3</sub>	$\begin{array}{c} \text{O} \\ \parallel \\ -\text{CCH}_2\text{CH}_3 \end{array}$		
a -120	CH <sub>3</sub>	$\begin{array}{c} \text{O} \\ \parallel \\ -\text{C}-\text{C}_3\text{H}_7(n) \end{array}$		m.p. 143 - 145°C
a -121	CH <sub>3</sub>	$\begin{array}{c} \text{O} \\ \parallel \\ -\text{C}-\text{C}_4\text{H}_9(n) \end{array}$		
a -122	CH <sub>3</sub>	$\begin{array}{c} \text{O} \\ \parallel \\ -\text{C}-\text{C}_6\text{H}_4-\text{CH}_3 \end{array}$		m.p. 179 - 182°C
a -123	CH <sub>3</sub>	$\begin{array}{c} \text{O} \\ \parallel \\ -\text{C}-\text{C}_6\text{H}_4-\text{Cl} \end{array}$		
a -124	CH <sub>3</sub>	-SO <sub>2</sub> -C <sub>6</sub> H <sub>4</sub> -CH <sub>3</sub>		
a -125	CH <sub>3</sub>	-SO <sub>2</sub> N(CH <sub>3</sub> ) <sub>2</sub>		m.p. 116 - 118°C
a -126	CH <sub>3</sub>	-SO <sub>2</sub> N(CH <sub>3</sub> ) <sub>2</sub>		m.p. 154 - 158°C

Table 4a (continued)
















Compound No.	R <sub>1</sub>	R <sub>2</sub>		Physical properties
a -127	CH <sub>3</sub>	-CON(CH <sub>3</sub> ) <sub>2</sub>		m.p. 136 - 138°C
a -128	CH <sub>3</sub>	-CON(CH <sub>3</sub> ) <sub>2</sub>		
a -129	CH <sub>3</sub>	-SO <sub>2</sub> N(CH <sub>3</sub> ) <sub>2</sub>		m.p. 50 - 60°C
a -130	CH <sub>3</sub>	-CON(CH <sub>3</sub> ) <sub>2</sub>		m.p. 200 - 203°C
a -131	CH <sub>3</sub>	-SO <sub>2</sub> - 		m.p. 57 - 60°C
a -132	CH <sub>3</sub>	-SO <sub>2</sub> - 		Viscous
a -133	CH <sub>3</sub>	-SO <sub>2</sub> - 		m.p. 105 - 107°C
a -134	CH <sub>3</sub>	-SO <sub>2</sub> - 		m.p. 131 - 133°C
a -135	CH <sub>3</sub>	-SO <sub>2</sub> - 		m.p. 169 - 172°C



Table 4a (continued)

















Compound No.	R <sub>1</sub>	R <sub>2</sub>		Physical properties
a -136	CH <sub>3</sub>			Oily
a -137	CH <sub>3</sub>			Viscous
a -138	CH <sub>3</sub>			m.p. 125 - 128°C
a -139	CH <sub>3</sub>	H		Oily
a -140	CH <sub>3</sub>			m.p. 139 - 142°C
a -141	CH <sub>3</sub>			m.p. 150 - 151°C
a -142	CH <sub>3</sub>	H		m.p. 95 - 103°C
a -143	CH <sub>3</sub>			m.p. 92 - 96°C
a -144	CH <sub>3</sub>	H		m.p. 60 - 70°C

Table 4a (continued)

















Compound No.	R <sub>1</sub>	R <sub>2</sub>	(X) <sub>n</sub>	Physical properties
a - 145	CH <sub>3</sub>	-SO <sub>2</sub> -  -CH <sub>3</sub>	SO <sub>2</sub> CH <sub>3</sub> - 	m.p. 75 - 80°C
a - 146	CH <sub>3</sub>	H	CF <sub>3</sub> -  -SCH <sub>3</sub>	m.p. 70 - 85°C
a - 147	CH <sub>3</sub>	-SO <sub>2</sub> -  -CH <sub>3</sub>	SO <sub>2</sub> CH <sub>3</sub> - 	
a - 148	CH <sub>3</sub>	-CHCO-    CH <sub>3</sub>	SO <sub>2</sub> CH <sub>3</sub> -  -CF <sub>3</sub>	m.p. 152 - 153°C
a - 149	CH <sub>3</sub>	-CH <sub>2</sub> CH=CH- 	SO <sub>2</sub> CH <sub>3</sub> -  -CF <sub>3</sub>	Viscous
a - 150	CH <sub>3</sub>	-CH <sub>2</sub> - 	SCH <sub>3</sub> - 	Viscous
a - 151	CH <sub>3</sub>	-CH <sub>2</sub> - 	SOCH <sub>3</sub> - 	Viscous
a - 152	CH <sub>3</sub>	-CH <sub>2</sub> - 	SO <sub>2</sub> CH <sub>3</sub> - 	m.p. 157 - 160°C
a - 153	CH <sub>3</sub>	-SO <sub>2</sub> C <sub>3</sub> H <sub>7</sub> (n)	SCH <sub>3</sub> - 	m.p. 82 - 85°C

Table 4a (continued)















Compound No.	R <sub>1</sub>	R <sub>2</sub>		Physical properties
a -154	CH <sub>3</sub>	-SO <sub>2</sub> C <sub>3</sub> H <sub>7</sub> (n)		Viscous
a -155	CH <sub>3</sub>	-SO <sub>2</sub> C <sub>3</sub> H <sub>7</sub> (n)		m.p. 165 - 169°C
a -156	CH <sub>3</sub>	H		Viscous
a -157	CH <sub>3</sub>			Viscous
a -158	CH <sub>3</sub>			m.p. 120 - 130°C
a -159	CH <sub>3</sub>	-CH <sub>2</sub> - 		Viscous
a -160	CH <sub>3</sub>	-CH <sub>2</sub> - 		m.p. 177 - 178°C
a -161	CH <sub>3</sub>	-CH <sub>2</sub> - 		m.p. 173 - 175°C

Table 4a (continued)
















Compound No.	R <sub>1</sub>	R <sub>2</sub>	(X) <sub>n</sub>	Physical properties
a - 162	CH <sub>3</sub>	-CH <sub>2</sub> - 		Viscous
a - 163	CH <sub>3</sub>	-CH <sub>2</sub> - 		Viscous
a - 164	CH <sub>3</sub>	-CH <sub>2</sub> - 		m.p. 65 - 75°C
a - 165	CH <sub>3</sub>	H		m.p. 208°C (decomposition)
a - 166	CH <sub>3</sub>	-SO <sub>2</sub> -  -CH <sub>3</sub>		m.p. 144 - 147°C
a - 167	CH <sub>3</sub>	-SO <sub>2</sub> -  -CH <sub>3</sub>		m.p. 120 - 141°C
a - 168	CH <sub>3</sub>	-CH <sub>2</sub> CO-C(CH <sub>3</sub> ) <sub>3</sub>		m.p. 140 - 144°C
a - 169	CH <sub>3</sub>	-CH <sub>2</sub> - 		m.p. 119 - 122°C
a - 170	CH <sub>3</sub>	-CH <sub>2</sub> C≡CH		Oily

Table 4a (continued)
















Compound No.	R <sub>1</sub>	R <sub>2</sub>		Physical properties
a -171	CH <sub>3</sub>	-CH <sub>2</sub> - 	OSO <sub>2</sub> CH <sub>3</sub> 	Viscous
a -172	CH <sub>3</sub>	H	CH <sub>2</sub> SCH <sub>3</sub> 	Oily
a -173	CH <sub>3</sub>	-CH <sub>2</sub> CN	SOCH <sub>3</sub> 	Viscous
a -174	CH <sub>3</sub>	-CH <sub>2</sub> CN	SO <sub>2</sub> CH <sub>3</sub> 	Viscous
a -175	CH <sub>3</sub>	-SO <sub>2</sub> -  -CH <sub>3</sub>	SCH <sub>3</sub> 	m.p. 153 - 156°C
a -176	CH <sub>3</sub>	-SO <sub>2</sub> -  -CH <sub>3</sub>	SO <sub>2</sub> CH <sub>3</sub> 	Viscous
a -177	CH <sub>3</sub>	-SO <sub>2</sub> -  -CH <sub>3</sub>	SCH <sub>3</sub> Cl 	Viscous
a -178	CH <sub>3</sub>	-SO <sub>2</sub> -  -CH <sub>3</sub>	SO <sub>2</sub> CH <sub>3</sub> Cl 	Viscous
a -179	CH <sub>3</sub>	-SO <sub>2</sub> C <sub>3</sub> H <sub>7</sub> (n)	SO <sub>2</sub> CH <sub>3</sub> Cl 	m.p. 140 - 144°C

Table 4a (continued)















Compound No.	R <sub>1</sub>	R <sub>2</sub>	(X) <sub>n</sub>	Physical properties
a - 180	CH <sub>3</sub>	-CH <sub>2</sub> - 	CH <sub>2</sub> SCH <sub>3</sub> 	Viscous
a - 181	CH <sub>3</sub>	-CH <sub>2</sub> - 	CH <sub>2</sub> SOCH <sub>3</sub> 	Viscous
a - 182	CH <sub>3</sub>	-CH <sub>2</sub> - 	CH <sub>2</sub> SO <sub>2</sub> CH <sub>3</sub> 	m.p. 55 - 62°C
a - 183	CH <sub>3</sub>	H	SCH <sub>3</sub> Cl 	Viscous
a - 184	CH <sub>3</sub>	-CH <sub>2</sub> - 	SCH <sub>3</sub> Cl 	Viscous
a - 185	CH <sub>3</sub>	-CH <sub>2</sub> - 	SOCH <sub>3</sub> Cl 	m.p. 154 - 155°C
a - 186	CH <sub>3</sub>	-CH <sub>2</sub> - 	SO <sub>2</sub> CH <sub>3</sub> Cl 	m.p. 165 - 167°C
a - 187	CH <sub>3</sub>	H	N(CH <sub>3</sub> )SO <sub>2</sub> CH <sub>3</sub> 	m.p. 50 - 58°C

Table 4a (continued)















Compound No.	R <sub>1</sub>	R <sub>2</sub>	(X) <sub>n</sub> 	Physical properties
a -188	CH <sub>3</sub>	-SO <sub>2</sub> C <sub>3</sub> H <sub>7</sub> (n)	N(CH <sub>3</sub> )SO <sub>2</sub> CH <sub>3</sub> 	m.p. 181 - 184°C
a -189	CH <sub>3</sub>	-SO <sub>2</sub> -  -CH <sub>3</sub>	N(CH <sub>3</sub> )SO <sub>2</sub> CH <sub>3</sub> 	m.p. 70 - 73°C
a -190	CH <sub>3</sub>	-CH <sub>2</sub> C≡CH	SCH <sub>3</sub> Cl 	Viscous
a -191	CH <sub>3</sub>	-CH <sub>2</sub> C≡CH	SO <sub>2</sub> CH <sub>3</sub> Cl 	m.p. 40 - 50°C
a -192	CH <sub>3</sub>	H	SCH <sub>3</sub> 	Viscous
a -193	CH <sub>3</sub>	-CH <sub>2</sub> - 	SCH <sub>3</sub> 	m.p. 107 - 110°C
a -194	CH <sub>3</sub>	-CH <sub>2</sub> - 	SOCH <sub>3</sub> 	m.p. 48 - 52°C
a -195	CH <sub>3</sub>	-CH <sub>2</sub> - 	SO <sub>2</sub> CH <sub>3</sub> 	m.p. 140 - 148°C
a -196	CH <sub>3</sub>	H	SCH <sub>3</sub> 	m.p. 115 - 130°C

Table 4a (continued)



















Compound No.	R <sub>1</sub>	R <sub>2</sub>		Physical properties
a -197	CH <sub>3</sub>			m.p. 79 - 93°C
a -198	CH <sub>3</sub>			m.p. 114 - 125°C
a -199	CH <sub>3</sub>			m.p. 143 - 146°C
a -200	CH <sub>3</sub>			m.p. 173 - 178°C
a -201	CH <sub>3</sub>	H		m.p. 155 - 156°C
a -202	CH <sub>3</sub>			m.p. 150 - 151°C
a -203	CH <sub>3</sub>			m.p. 183 - 188°C
a -204	CH <sub>3</sub>			Viscous
a -205	CH <sub>3</sub>			m.p. 111 - 126°C



Table 4a (continued)












Compound No.	R <sub>1</sub>	R <sub>2</sub>	(X) <sub>n</sub> 	Physical properties
a -206	CH <sub>3</sub>	-CH <sub>2</sub> C≡CH		Oily
a -207	CH <sub>3</sub>	-SO <sub>2</sub> C <sub>3</sub> H <sub>7</sub> (n)		Viscous
a -208	CH <sub>3</sub>	-SO <sub>2</sub> C <sub>3</sub> H <sub>7</sub> (n)		m.p. 134 - 135°C
a -209	CH <sub>3</sub>	-CH <sub>2</sub> C≡CH		Viscous
a -210	CH <sub>3</sub>	-CH <sub>2</sub> - 		Viscous
a -211	CH <sub>3</sub>	H		m.p. 122 - 138°C
a -212	CH <sub>3</sub>	-CH <sub>2</sub> CN		Viscous
a -213	CH <sub>3</sub>	-CH <sub>2</sub> C=CH <sub>2</sub>   Cl		m.p. 108 - 111°C
a -214	CH <sub>3</sub>	-SO <sub>2</sub> (CH <sub>2</sub> ) <sub>3</sub> Cl		m.p. 113 - 118°C

Table 4a (continued)












Compound No.	R <sub>1</sub>	R <sub>2</sub>	(X) <sub>n</sub>	Physical properties
a -215	CH <sub>3</sub>	-CH <sub>2</sub> - 		m.p. 50 - 60°C
a -216	CH <sub>3</sub>	-CO(CH <sub>2</sub> ) <sub>3</sub> Cl		m.p. 133 - 135°C
a -217	CH <sub>3</sub>	-CH <sub>2</sub> CH=CH <sub>2</sub>		Viscous
a -218	CH <sub>3</sub>	-CH <sub>2</sub> CN		Viscous
a -219	CH <sub>3</sub>	-CH <sub>2</sub> - 		Viscous
a -220	CH <sub>3</sub>	-SO <sub>2</sub> CH <sub>3</sub>		m.p. 125 - 130°C
a -221	CH <sub>3</sub>	-CO <sub>2</sub> CH <sub>3</sub>		Viscous
a -222	CH <sub>3</sub>	H		m.p. 153 - 156°C
a -223	CH <sub>3</sub>	-CH <sub>2</sub> CO <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>		Viscous

Table 4a (continued)












Compound No.	R <sub>1</sub>	R <sub>2</sub>		Physical properties
a -224	CH <sub>3</sub>	H		
a -225	CH <sub>3</sub>	H		Viscous
a -226	CH <sub>3</sub>	-SO <sub>2</sub> C <sub>3</sub> H <sub>7</sub> (n)		Viscous
a -227	CH <sub>3</sub>	H		Oily
a -228	CH <sub>3</sub>	-CH <sub>2</sub> CH=CHCl		m.p. 100 - 103°C
a -229	CH <sub>3</sub>	$\begin{array}{c} -\text{CH}_2\text{C}=\text{CH}_2 \\   \\ \text{CH}_3 \end{array}$		m.p. 122 - 125°C
a -230	CH <sub>3</sub>	$\begin{array}{c} -\text{CH}_2\text{C}=\text{CH}_2 \\   \\ \text{Br} \end{array}$		Oily
a -231	CH <sub>3</sub>	-CH <sub>2</sub> CH=C(CH <sub>3</sub> ) <sub>2</sub>		
a -232	CH <sub>3</sub>	-CH <sub>2</sub> CH=C(Cl) <sub>2</sub>		m.p. 142 - 145°C
a -233	CH <sub>3</sub>	-CH <sub>2</sub> CH=CHCH <sub>3</sub>		m.p. 101 - 106°C

Table 4a (continued)












Compound No.	R <sub>1</sub>	R <sub>2</sub>		Physical properties
a - 234	CH <sub>3</sub>	$\begin{array}{c} -\text{CHCH}=\text{CH}_2 \\   \\ \text{CH}_3 \end{array}$		Oily
a - 235	CH <sub>3</sub>	$-\text{CH}_2\text{CON}(\text{CH}_3)_2$		
a - 236	CH <sub>2</sub> CH <sub>3</sub>	$\begin{array}{c} -\text{CH}_2\text{C}=\text{CH}_2 \\   \\ \text{Cl} \end{array}$		
a - 237	CH <sub>2</sub> CH <sub>3</sub>	$\begin{array}{c} -\text{CH}_2\text{C}=\text{CH}_2 \\   \\ \text{CH}_3 \end{array}$		
a - 238	CH <sub>3</sub>	$\begin{array}{c} -\text{CH}_2\text{C}=\text{CH}_2 \\   \\ \text{CH}_3 \end{array}$		
a - 239	CH <sub>3</sub>	$\begin{array}{c} -\text{CH}_2\text{C}=\text{CH}_2 \\   \\ \text{Br} \end{array}$		m.p. 124 - 126°C
a - 240	CH <sub>3</sub>	$-\text{CH}_2\text{CH}=\text{C}(\text{CH}_3)_2$		
a - 241	CH <sub>3</sub>	$\begin{array}{c} -\text{CHCH}=\text{CH}_2 \\   \\ \text{CH}_3 \end{array}$		Viscous
a - 242	CH <sub>3</sub>	$-\text{SO}_2\text{CH}_2\text{CH}_3$		
a - 243	CH <sub>3</sub>	$-\text{SO}_2\text{C}_6\text{H}_5(n)$		

Table 4a (continued)












Compound No.	R <sub>1</sub>	R <sub>2</sub>		Physical properties
a -244	CH <sub>3</sub>	H		m.p. 60 - 65°C
a -245	CH <sub>3</sub>	-SO <sub>2</sub> C <sub>3</sub> H <sub>7</sub> (n)		
a -246	CH <sub>3</sub>	$\begin{array}{c} -\text{CH}_2\text{C}=\text{CH}_2 \\   \\ \text{Cl} \end{array}$		
a -247	CH <sub>3</sub>	$\begin{array}{c} -\text{CH}_2\text{C}=\text{CH}_2 \\   \\ \text{Cl} \end{array}$		
a -248	CH <sub>3</sub>	$\begin{array}{c} -\text{CH}_2\text{C}=\text{C}(\text{CH}_3)_2 \\   \\ \text{CH}_3 \end{array}$		
a -249	CH <sub>3</sub>	-CH <sub>2</sub> CH=CH <sub>2</sub>		Viscous
a -250	CH <sub>3</sub>	-CH <sub>2</sub> CN		Viscous
a -251	CH <sub>3</sub>	$\begin{array}{c} -\text{CH}_2\text{C}=\text{CH}_2 \\   \\ \text{Cl} \end{array}$		Viscous
a -252	CH <sub>3</sub>	-CH <sub>2</sub> CH=CHCl		Viscous
a -253	CH <sub>3</sub>	-CH <sub>2</sub> CH=CHCl		Viscous

Table 4a (continued)





















Compound No.	R <sub>1</sub>	R <sub>2</sub>		Physical properties
a -254	CH <sub>3</sub>	$-\text{CH}_2\text{C}(\text{Cl})=\text{CH}_2$		m.p. 120 - 125°C
a -255	CH <sub>3</sub>	$-\text{CH}_2\text{CH}=\text{CHCH}_3$		m.p. 97 - 105°C
a -256	CH <sub>3</sub>	$-\text{CH}_2\text{CH}=\text{CHCH}_3$		m.p. 116 - 117°C
a -257	CH <sub>3</sub>	$-\text{SO}_2\text{CH}_2\text{CH}_3$		Viscous
a -258	CH <sub>3</sub>	$-\text{SO}_2\text{CH}_3$		m.p. 164 - 165°C
a -259	CH <sub>3</sub>	$-\text{CH}_2\text{CO}_2\text{CH}_2\text{CH}_3$		Viscous
a -260	CH <sub>3</sub>	$-\text{SO}_2\text{CH}_2\text{CH}_3$		m.p. 112 - 115°C
a -261	CH <sub>3</sub>	$-\text{SO}_2\text{CH}_2\text{CH}_3$		Viscous
a -262	CH <sub>3</sub>	$-\text{CH}_2\text{C}(=\text{O})\text{CH}_2\text{C}(=\text{O})\text{CH}_2\text{CH}_3$		Viscous
a -263	CH <sub>3</sub>	$-\text{CH}_2\text{CON}(\text{C}_2\text{H}_5)_2$		Viscous

Table 4 (continued)

Compound No.	R <sub>1</sub>	R <sub>2</sub>		Physical properties
a -264	CH <sub>3</sub>	$-\text{CH}_2\text{CH}(\text{CH}_3)\text{CH}_2\text{Cl}$		m.p. 123~125°C
a -265	CH <sub>3</sub>	$-\text{SO}_2\text{C}_3\text{H}_7 (n)$		
a -266	CH <sub>3</sub>	$-\text{SO}_2\text{C}_3\text{H}_7 (n)$		
a -267	CH <sub>3</sub>	$-\text{CH}_2\text{CH}_2\text{Cl}$		
a -268	CH <sub>3</sub>	$-\text{CH}_2\text{CH}_2\text{Cl}$		
a -269	CH <sub>3</sub>	$-\text{COC}(\text{CH}_3)=\text{CH}_2$		
a -270	CH <sub>3</sub>	$-\text{SO}_2-\text{C}_6\text{H}_4-\text{CH}_3$		m.p. 67~72°C
a -271	CH <sub>3</sub>	$-\text{CH}_2\text{CH}=\text{C}(\text{Cl})_2$		Viscous




Table 4b

Comp. No.	R <sub>1</sub>	(Z)	R <sub>2</sub>	(X) <sub>n</sub>	Physical properties
b-1	CH <sub>3</sub>	CH <sub>3</sub> ~~~~~	-SO <sub>2</sub> --CH <sub>3</sub>	-SO <sub>2</sub> -CH <sub>3</sub> -CF <sub>3</sub>	m.p. 46-56°C
b-2	CH <sub>3</sub>	CH <sub>3</sub> ~~~~~	-CH <sub>2</sub> -	-SO <sub>2</sub> -CH <sub>3</sub> -CF <sub>3</sub>	Viscous
b-3	CH <sub>3</sub>	CH <sub>3</sub> ~~~~~		-SO <sub>2</sub> -CH <sub>3</sub> -CF <sub>3</sub>	
b-4	CH <sub>3</sub>	CH <sub>3</sub> ~~~~~	-SO <sub>2</sub> C <sub>3</sub> H <sub>7</sub> (n)	-SO <sub>2</sub> -CH <sub>3</sub> -CF <sub>3</sub>	
b-5	CH <sub>3</sub>	CH <sub>3</sub> ~~~~~	-CH <sub>2</sub> -	-SO <sub>2</sub> -CH <sub>3</sub> -CF <sub>3</sub>	
b-6	CH <sub>3</sub>	CH <sub>3</sub> ~~~~~	H	-SO <sub>2</sub> -CH <sub>3</sub> -CF <sub>3</sub>	



Now, the Test Examples of the present invention will be described.

#### TEST EXAMPLE 1

Upland field soil was put into a 1/150,000ha pot, and  
5 seeds of various plants were sown. Then, when the plants  
reached predetermined leaf stages (① barnyardgrass  
(Echinochloa crus-galli L.), EC: 1.3-2.6 leaf stage, ②  
crabgrass (Digitaria sanguinalis L.), DS: 1.0-2.5 leaf  
stage, ③ redroot pigweed (Amaranthus retroflexus L.),  
10 AR: 0.1-1.2 leaf stage, ④ prickly sida (Sida spinosa  
L.), SS: 0.1-1.2 leaf stage, ⑤ tall morningglory  
(Pharbitis purpurea L.), PP: 0.3-1.3 leaf stage, ⑥  
common cocklebur (Xanthium strumarium L.), XS: 0.1-1.8  
leaf stage, ⑦ rice (Oryza sativa L.), OS: 1.0-2.5 leaf  
15 stage, ⑧ wheat (Triticum spp.), TR: 2.2-2.9 leaf stage,  
⑨ corn (Zea mays L.), ZM: 1.8-3.5 leaf stage, ⑩ soybean  
(Glycine max Merr.), GM: primary leaf - 0.3 leaf stage),  
a wettable powder having the compound of the present  
invention formulated in accordance with a usual  
20 formulation method, was weighed so that the active  
ingredient would be a predetermined amount, and diluted  
with water in an amount of 500 g/ha. To the diluted  
solution, 0.1 % (v/v) of an agricultural spreader was  
added. The herbicide thus adjusted was applied by a  
25 small size spray for foliage treatment. On the 18th to  
30th days after the application of the herbicide, the  
growth of the respective plants was visually observed,

and the herbicidal effects were evaluated by the growth-controlling degrees (%) ranging from 0 (equivalent to the non-treated area) to 100 (complete kill), whereby the results shown in Table 5, were obtained. Compound Nos. 5 in Table 5 correspond to Compound Nos. in Table 4a and 4b given hereinbefore.

Table 5

Comp. No.	Dose of active ingredient (g/ha)	Growth-controlling degree (%)										Evaluation day
		EC	DS	AR	SS	PP	XS	OS	TR	ZM	GM	
a-3	500	0	0	70	60	60	70	40	0	0	0	23
a-5	500	30	40	70	0	60	60	50	10	0	10	18
a-11	63	40	20	90	20	80	80	50	0	0	50	22
	125	70	40	90	20	80	100	80	0	0	70	
	500	90	50	100	50	100	100	100	0	0	80	
a-12	63	80	40	90	30	80	95	80	0	20	70	18
	125	80	70	80	20	90	80	90	0	0	70	
	500	90	80	90	60	90	80	100	0	0	80	
a-13	500	80	60	90	60	80	100	90	0	0	60	18
a-24	500	40	50	80	10	60	80	50	-	10	30	18
a-25	500	70	90	80	10	70	90	50	-	30	60	18
a-27	500	60	20	100	20	80	80	50	0	0	40	20
a-39	63	70	30	80	20	60	80	50	0	10	60	20
	125	80	60	90	30	60	100	70	0	10	60	
	500	90	80	90	30	70	100	70	10	20	80	
a-40	63	90	30	90	10	90	80	90	-	30	50	18
	125	90	40	90	30	90	90	100	-	50	70	
	500	100	90	100	60	90	100	100	-	70	70	
a-41	63	90	20	90	20	90	70	90	-	0	50	18
	125	90	40	90	20	90	80	90	-	0	70	
	500	100	90	100	40	100	100	100	-	20	90	
a-42	500	60	40	100	30	70	80	50	10	20	50	20
a-43	125	50	30	90	20	60	80	40	0	20	50	20
	500	70	80	100	30	80	100	70	20	20	70	
a-44	63	80	50	80	20	60	80	60	-	0	60	18
	125	90	70	80	20	70	90	80	-	20	70	
	500	90	90	90	30	90	100	80	-	20	80	
a-45	125	40	30	80	20	80	60	10	10	40	50	20
	500	70	50	90	20	80	100	20	20	20	50	
a-47	125	40	40	90	30	60	80	20	-	0	0	18
	500	80	50	100	60	80	80	70	-	30	40	

Table 5. (continued)

Comp. No.	Dose of active ingredient (g/ha)	Growth-controlling degree (%)										Evaluation day
		EC	DS	AR	SS	PP	XS	OS	TR	ZM	GM	
a-48	63	80	40	90	10	40	70	50	-	0	50	18
	125	90	60	90	30	70	100	60	-	20	60	
	500	100	80	100	60	90	100	80	-	40	70	
a-49	125	80	40	100	30	70	90	60	10	20	60	20
	500	80	70	100	30	80	90	80	20	20	60	
a-51	125	90	60	80	10	80	80	70	-	0	70	19
	500	90	90	100	30	90	90	80	-	10	70	
a-52	63	90	60	80	20	70	80	60	-	20	80	18
	125	90	70	80	30	80	90	70	-	30	90	
	500	90	90	90	50	90	100	80	-	40	90	
a-53	63	70	10	90	0	70	80	50	-	0	50	18
	125	80	20	90	10	80	90	50	-	0	60	
	500	90	90	90	20	90	100	90	-	10	70	
a-59	500	70	40	80	0	80	90	30	0	10	60	20
a-60	63	90	50	80	10	90	90	60	-	0	100	19
	125	90	80	90	20	100	100	70	-	20	90	
	500	90	90	100	30	100	100	90	-	60	90	
a-69	500	40	40	90	50	70	100	60	0	10	40	20
a-72	125	70	20	80	0	60	90	50	-	0	40	19
	500	100	30	90	0	70	100	80	-	0	60	
a-73	500	60	30	90	40	70	80	10	-	30	50	18
a-76	63	70	30	90	0	80	90	70	0	10	70	20
	125	80	60	90	20	80	90	70	10	20	70	
	500	90	90	100	40	100	100	70	20	40	80	
a-77	125	90	30	80	10	70	70	50	-	10	60	18
	500	90	60	90	30	70	90	90	-	30	70	
a-78	63	80	40	70	20	60	80	40	-	10	60	18
	125	80	40	90	20	90	90	60	-	20	70	
	500	100	50	100	40	90	100	60	-	50	70	
a-79	500	60	30	100	50	80	90	30	-	0	70	18
a-80	500	80	90	80	20	60	80	80	-	60	10	19

Table 5 (continued)

Comp. No.	Dose of active ingredient (g/ha)	Growth-controlling degree (%)										Evaluation day
		EC	DS	AR	SS	PP	XS	OS	TR	ZM	GM	
a-81	125	60	60	90	10	90	40	10	-	0	30	19
	500	90	90	90	30	90	-	40	-	10	50	
a-82	125	40	30	80	20	60	100	60	-	0	60	19
	500	80	30	90	20	90	100	80	-	10	70	
a-83	63	50	60	80	30	70	100	60	-	0	10	18
	125	60	70	80	40	80	100	60	-	10	20	
	500	90	90	90	40	90	90	80	-	30	70	
a-84	63	70	30	70	30	40	100	30	-	0	50	18
	125	80	50	80	20	60	100	60	-	0	60	
	500	90	90	90	50	90	90	100	-	20	70	
a-87	125	80	20	90	10	70	100	60	-	10	60	19
	500	100	70	95	10	100	-	80	-	0	95	
a-88	63	90	60	80	30	80	90	70	-	10	70	18
	125	90	80	90	60	80	90	70	-	20	80	
	500	100	90	90	60	90	100	80	-	60	90	
a-89	63	90	90	90	30	90	100	100	-	10	90	19
	125	90	90	90	30	90	100	90	-	20	100	
	500	100	100	100	60	100	100	100	-	80	100	
a-90	125	80	90	70	0	60	80	60	-	40	10	19
	500	90	100	80	10	70	100	80	-	80	70	
a-92	500	80	90	90	20	90	-	50	-	0	20	19
a-93	125	60	40	95	0	60	100	10	-	0	70	19
	500	80	50	100	20	100	-	70	-	0	100	
a-94	63	90	90	80	10	90	80	70	-	0	60	19
	125	100	90	90	-	100	100	90	-	10	70	
	500	90	90	90	20	100	100	90	-	20	70	
a-95	63	100	70	100	10	80	90	90	-	20	95	19
	125	100	90	100	10	95	100	100	-	35	95	
	500	100	100	100	40	100	-	100	-	70	100	
a-96	125	80	70	80	20	80	90	60	10	40	80	20
	500	100	100	100	30	100	100	70	20	60	90	
a-97	63	90	30	90	0	40	50	10	-	0	20	18
	125	80	40	80	10	50	70	30	-	0	30	
	500	90	90	90	20	70	80	50	-	20	60	

Table 5 (continued)

Comp. No.	Dose of active ingredient (g/ha)	Growth-controlling degree (%)										Evaluation day
		EC	DS	AR	SS	PP	XS	OS	TR	ZM	GM	
a-98	63	90	30	90	0	60	70	30	-	0	40	18
	125	90	40	100	30	80	70	40	-	0	50	
	500	90	90	90	30	90	100	100	-	20	60	
a-99	63	90	40	100	40	80	80	70	-	40	70	18
	125	100	60	100	40	90	100	90	-	50	70	
	500	100	100	100	70	100	100	100	-	80	100	
a-100	125	80	40	90	30	80	70	0	-	0	50	18
	500	90	90	100	40	90	100	50	-	10	60	
a-101	125	90	30	90	0	30	20	0	-	0	50	18
	500	90	40	90	30	30	60	0	-	20	60	
a-102	63	90	40	90	20	70	80	80	-	0	100	19
	125	95	60	90	30	80	80	100	-	10	60	
	500	100	90	100	70	90	80	90	-	20	100	
a-103	63	90	30	90	40	60	70	50	-	0	50	19
	125	100	50	60	40	80	100	50	-	0	50	
	500	100	90	80	50	100	100	100	-	40	70	
a-104	125	80	20	90	0	10	0	0	-	0	10	19
	500	90	20	100	10	10	0	0	-	0	20	
a-105	63	90	30	100	30	40	20	10	-	10	40	19
	125	90	40	100	0	60	80	30	-	0	50	
	500	100	80	80	30	90	-	50	-	80	90	
a-106	63	10	30	80	50	60	-	20	-	0	30	21
	125	70	70	100	50	90	90	90	-	20	40	
	500	90	70	70	-	100	50	100	-	50	80	
a-116	125	50	10	100	30	70	-	30	-	10	30	24
	500	50	50	90	20	90	-	20	-	0	40	
a-118	63	70	30	100	30	70	90	100	-	10	70	21
	125	80	50	80	30	80	-	100	-	40	100	
	500	90	80	100	-	100	100	100	-	60	100	
a-120	63	90	50	100	20	50	80	90	-	0	70	21
	125	100	50	100	50	80	100	100	-	0	80	
	500	90	90	-	100	60	100	100	-	70	100	
a-122	63	70	50	100	40	60	90	60	-	10	50	21
	125	80	90	100	50	70	90	90	-	30	60	
	500	90	90	-	-	50	100	100	-	40	70	

Table 5 (continued)

Comp. NO.	Dose of active ingredient (g/ha)	Growth-controlling degree (%)										Evaluation day
		EC	DS	AR	SS	PP	XS	OS	TR	ZM	GM	
a-131	500	80	40	60	30	80	100	50	-	30	20	19
a-132	125 500	80 90	70 90	60 60	20 20	90 90	50 50	10 40	- -	30 70	0 10	19
a-133	500	10	30	90	40	70	10	0	-	40	30	21
a-136	125 500	20 100	30 100	90 80	50 30	70 70	- 50	0 0	- -	40 100	40 50	21
a-137	125 500	20 90	30 60	90 100	0 10	100 100	- 80	0 40	- -	40 80	30 50	21
a-138	500	100	80	90	70	100	20	0	-	90	50	21
a-139	500	10	50	90	50	70	70	0	-	80	50	21
a-140	63 125 500	90 90 100	10 30 90	100 100 -	10 20 30	40 80 100	70 80 80	0 20 90	- - -	0 10 0	30 40 50	21
a-141	125 500	80 90	60 90	100 -	30 60	90 100	90 100	90 90	- -	20 30	70 80	21
a-143	500	90	30	50	20	80	100	20	-	20	60	18
a-144	500	50	90	30	40	60	100	50	-	0	20	18
a-146	125 500	20 60	30 80	80 80	40 40	80 90	50 100	40 60	- -	20 30	40 70	19
a-148	125 500	60 90	10 80	100 100	50 50	70 80	10 20	20 20	- -	10 40	10 40	21
a-149	63 125 500	60 90 100	20 30 80	90 80 100	40 50 70	80 80 100	- - 100	0 40 100	- - -	10 20 60	60 70 90	21
a-150	125 500	90 100	70 90	50 70	20 20	80 100	- 70	40 70	- -	100 100	60 70	25

Table 5 (continued)

Comp. No.	Dose of active ingredient (g/ha)	Growth-controlling degree (%)										Evaluation day
		EC	DS	AR	SS	PP	XS	OS	TR	ZM	GM	
a-151	125 500	100 100	90 90	40 100	30 30	80 100	30 80	20 20	- -	50 90	30 50	25
a-152	125 500	100 100	60 90	70 90	10 50	60 100	20 100	10 20	- -	60 50	10 40	25
a-153	125 500	60 90	20 60	50 60	10 30	40 50	80 -	10 20	- -	0 10	0 20	24
a-154	500	90	70	70	10	70	60	50	-	0	30	25
a-155	500	60	70	100	10	70	100	20	-	0	40	25
a-156	250	90	70	20	20	70	-	20	-	10	40	21
a-158	125 250	60 70	40 50	90 90	20 20	70 90	70 -	50 50	- -	40 60	50 50	21
a-162	500	60	20	0	10	80	-	10	0	10	0	21
a-164	500	70	70	20	10	90	-	10	0	40	50	21
a-165	63 125 500	100 100 100	95 95 100	100 100 100	40 100 100	100 100 100	50 - -	99 100 100	50 50 80	50 90 95	40 60 100	30
a-166	63 125 500	95 100 100	80 95 100	100 - 100	40 50 50	90 100 100	20 20 -	70 100 100	10 50 50	40 40 70	40 40 50	30
a-167	125 500	10 40	20 30	100 100	0 90	100 100	90 100	40 70	- -	0 30	40 100	21
a-168	63 125 500	90 100 100	60 70 100	100 100 100	50 60 60	90 100 100	90 - -	100 100 100	- - -	40 70 100	50 70 100	21
a-169	63 125	90 90	60 70	100 90	30 40	100 100	10 10	10 20	- -	10 10	30 60	21
a-170	63 125	90 90	60 80	80 100	20 50	100 100	70 70	90 100	- -	10 60	90 100	21



Table 5 (continued)

Comp. NO.	Dose of active ingredient (g/ha)	Growth-controlling degree (%)										Evaluation day
		EC	DS	AR	SS	PP	XS	OS	TR	ZM	GM	
a-171	500	40	10	40	0	80	-	0	0	0	60	21
a-172	125 500	60 100	20 50	80 90	30 60	80 100	80 -	80 100	0 0	0 50	50 95	21
a-173	125 500	70 90	30 60	30 40	10 30	80 100	30 -	100 100	0 0	0 40	50 60	21
a-174	125 500	70 100	40 60	50 90	0 30	80 100	70 -	100 80	0 0	0 50	70 80	21
a-177	500	80	50	30	0	90	-	10	0	20	40	21
a-178	500	60	40	60	0	80	-	50	0	0	50	21
a-179	500	70	70	90	90	70	-	50	0	0	80	21
a-180	125 500	80 100	20 50	50 90	20 30	100 100	50 -	100 50	0 0	0 60	70 80	21
a-181	125 500	70 100	20 40	70 60	10 20	100 100	70 -	50 90	0 0	0 40	70 80	21
a-182	125 500	60 60	20 60	80 80	0 10	100 100	60 -	50 50	0 0	0 20	50 90	21
a-183	500	0	40	10	10	100	100	0	0	10	40	21
a-184	500	50	20	10	20	100	-	0	0	0	40	21
a-187	500	70	20	100	40	95	-	50	0	0	60	21
a-188	500	50	10	90	40	95	-	10	0	0	70	21
a-189	500	90	10	100	40	95	-	90	0	0	70	21
a-191	125 500	90 100	30 50	90 90	20 40	90 90	- -	20 40	0 0	0 0	50 50	21

Table 5 (continued)

Comp. No.	Dose of active ingredient (g/ha)	Growth-controlling degree (%)										Evaluation day
		EC	DS	AR	SS	PP	XS	OS	TR	ZM	GM	
a-192	500	30	0	30	0	80	90	30	0	10	60	21
a-194	500	0	30	20	0	100	90	0	0	0	50	22
a-200	500	70	50	95	20	100	90	20	0	0	40	26
a-201	500	20	40	90	40	20	70	40	0	0	0	26
a-203	125 500	20 70	60 80	90 100	10 50	80 100	80 95	20 90	0 0	0 0	60 90	22
a-204	500	70	50	30	0	80	80	20	0	0	50	22
a-206	125 500	80 100	30 80	30 100	20 50	100 100	100 100	90 100	0 0	0 70	90 100	22
a-207	125 500	70 90	20 60	20 90	0 40	100 100	80 100	80 100	0 0	0 20	80 100	22
a-208	125 500	30 70	40 60	90 70	10 40	80 90	100 100	50 95	0 0	0 0	50 60	22
a-209	500	40	10	30	0	90	80	70	0	0	70	22
a-210	125 500	90 100	60 90	70 100	10 40	90 100	90 90	90 90	0 0	0 60	80 100	27
a-211	125 500	40 80	20 50	70 90	0 20	50 70	80 90	70 70	0 0	0 0	40 50	27
a-212	125 500	70 100	0 10	40 70	0 0	0 90	70 80	60 80	0 0	0 0	50 80	27
a-213	125 500	100 100	80 100	95 100	30 70	100 100	100 100	80 100	0 0	20 70	95 100	26
a-214	125 500	60 100	60 90	100 100	20 70	90 100	70 90	100 100	0 0	0 0	50 70	27
a-215	500	0	0	90	20	90	80	0	0	0	40	23

Table 5 (continued)

Comp. No.	Dose of active ingredient (g/ha)	Growth-controlling degree (%)										Evaluation day
		EC	DS	AR	SS	PP	XS	OS	TR	ZM	GM	
a-216	125 500	50 90	70 90	90 100	10 60	70 80	70 100	50 100	0 0	0 0	40 70	23
a-217	125 500	60 30	30 60	50 80	10 20	100 100	80 100	70 100	0 0	0 70	70 100	20
a-218	125 500	70 90	20 60	10 50	10 20	90 100	70 80	70 90	0 0	0 40	80 100	20
a-219	125 500	70 100	30 70	10 60	0 30	100 100	80 100	50 80	0 0	0 70	80 100	20
a-220	125 500	10 70	10 30	10 30	0 20	70 90	80 80	10 60	0 0	0 0	40 40	20
a-221	125 500	60 80	0 60	0 40	0 30	80 100	80 100	50 50	0 0	0 0	70 90	20
a-222	125 500	30 80	10 50	30 80	10 50	80 90	70 90	20 70	0 0	0 10	40 60	20
a-257	125 500	70 90	10 60	10 10	0 10	90 100	70 100	20 30	0 0	0 30	50 100	28
a-258	125 500	50 90	10 70	30 80	0 30	90 100	70 100	10 50	0 0	0 0	40 70	28
a-259	125 500	70 90	70 100	0 20	0 0	80 90	90 100	10 40	0 0	0 0	60 80	28
b-1	500	80	50	80	20	90	-	60	-	20	60	19
b-2	500	90	20	70	20	90	60	60	-	0	60	18

## TEST EXAMPLE 2

Paddy field soil was put into a 1/1,000,000ha pot, and seeds of barnyardgrass (Echinochloa crus-galli L.) and japanese bulrush (Scirpus juncoides) were sown and  
5 slightly covered with soil. Then, the pot was left to stand still in a greenhouse in a state where the depth of flooding water was from 0.5 to 1 cm, and two days later, tubers of japanese ribbon wapato (Sagittaria pygmaea) were planted. Thereafter, the depth of flooding water  
10 was maintained at a level of from 3 to 4 cm, and when barnyardgrass and japanese bulrush reached a 0.5 leaf stage and japanese ribbon wapato reached to a primary leaf stage, an aqueous diluted solution of a wettable powder having the compound of the present invention  
15 formulated in accordance with a usual formulation method, was uniformly applied under submerged condition by a pipette so that the dose of the active ingredient would be at a predetermined level.

On the other hand, paddy field soil was put into a  
20 1/1,000,000ha pot and puddled and leveled, and the depth of flooding water was from 3 to 4 cm. One day later, rice (Oryza sativa L. var. Nihonbare) of 2 leaf stage was transplanted in a depth of 3 cm. On the 4th day after the transplantation, the compound of the present  
25 invention was applied in the same manner as described above.

On the 14th days after the application of the

herbicide, the growth of barnyard grass, japanese  
burlrush and japanese ribbon wapato was visually observed  
and on the 21st day after the application the herbicide,  
the growth of rice was visually observed, and the  
5 herbicidal effects were evaluated by the growth-  
controlling degrees (%) ranging from 0 (equivalent to the  
non-treated area) to 100 (complete kill), whereby the  
results shown in Table 6 were obtained. Compound Nos. in  
Table 6 correspond to Compound Nos. in Table 4a and 4b  
10 given hereinbefore. The growth controlling degrees  
against rice of compounds Nos. a-101 et seq (except for  
a-131, a-132, a-145, a-146 and b-1) are mean values of  
two test results.

Table 6

Comp. No.	Dose of active ingredient (g/ha)	Growth-controlling degree (%)			
		EC	SJ	SP	OS
a-1	1 0 0 0 5 0 0	5 0 4 0	1 0 0 1 0 0	1 0 0 —	— —
a-2	5 0 0 2 5 0	9 0 8 5	9 5 8 5	9 0 8 5	0 0
a-3	1 0 0 0 5 0 0	9 5 9 5	1 0 0 1 0 0	9 5 8 5	1 0 0
a-4	5 0 0 2 5 0	1 0 0 1 0 0	8 5 5 0	8 5 6 0	0 0
a-5	1 0 0 0 5 0 0	4 0 0	8 5 8 5	8 5 7 0	9 0 3 0
a-6	5 0 0 2 5 0	1 0 0 1 0 0	9 0 8 5	9 0 8 5	5 0 3 0
a-7	5 0 0 2 5 0	1 0 0 9 9	1 0 0 9 5	8 5 8 5	8 0 7 0
a-8	1 0 0 0 5 0 0	0 0	8 5 5 0	8 5 8 5	0 0
a-9	5 0 0	1 0	5 0	7 0	3 0
a-10	5 0 0 2 5 0	8 0 6 0	5 0 5 0	8 5 8 5	0 0
a-11	2 5 0 1 2 5	1 0 0 1 0 0	9 5 9 0	9 0 8 5	1 0 0 1 0 0
a-12	5 0 0 2 5 0	1 0 0 1 0 0	1 0 0 1 0 0	9 0 9 0	1 0 0 1 0 0
a-13	5 0 0 2 5 0	1 0 0 1 0 0	1 0 0 1 0 0	9 0 9 0	1 0 0 1 0 0
a-14	5 0 0 2 5 0	4 0 1 0	8 5 8 5	9 0 8 5	1 0 0
a-15	5 0 0 2 5 0	1 0 0 5 0	8 5 7 0	7 0 7 0	0 0
a-16	5 0 0 2 5 0	1 0 0 9 9	5 0 0	3 0 3 0	3 0 0
a-17	5 0 0 2 5 0	1 0 0 9 9	8 5 5 0	5 0 5 0	2 0 0

Table 6 (continued)

Comp. No.	Dose of active ingredient (g/ha)	Growth-controlling degree (%)			
		EC	SJ	SP	OS
a-18	5 0 0	0	9 0	9 5	0
	2 5 0	0	8 5	9 5	0
a-19	5 0 0	7 0	9 0	9 0	1 0
	2 5 0	4 0	9 0	8 5	0
a-20	5 0 0	9 0	8 5	3 0	6 0
	2 5 0	8 5	5 0	0	3 5
a-21	5 0 0	5 0	9 5	9 0	3 0
	2 5 0	3 0	9 0	5 0	3 0
a-22	5 0 0	9 5	9 5	9 0	3 0
	2 5 0	1 0 0	9 5	5 0	2 0
a-23	5 0 0	1 0 0	9 9	9 0	0
	2 5 0	5 0	9 9	5 0	0
a-24	2 5 0	4 0	8 5	9 0	1 0
	1 2 5	2 0	8 0	8 5	0
a-25	2 5 0	1 0 0	9 0	9 0	0
	1 2 5	8 5	8 5	8 5	0
a-26	1 0 0 0	9 0	9 0	1 0	1 0
	5 0 0	5 0	9 0	1 0	0
a-27	2 5 0	1 0 0	9 5	9 0	1 0 0
	1 2 5	1 0 0	9 0	8 5	4 0
a-39	1 2 5	1 0 0	9 0	8 5	3 5
	6 3	1 0 0	8 5	5 0	3 0
a-40	1 2 5	1 0 0	1 0 0	9 0	8 0
	6 3	1 0 0	9 9	9 0	4 0
a-41	1 2 5	1 0 0	9 5	8 5	1 0 0
	6 3	1 0 0	9 0	8 5	7 0
a-42	2 5 0	1 0 0	9 0	9 0	9 0
	1 2 5	1 0 0	9 0	0	9 0
a-43	2 5 0	1 0 0	9 0	9 0	9 0
	1 2 5	1 0 0	8 5	5 0	9 9
a-44	2 5 0	1 0 0	9 5	9 0	9 5
	1 2 5	1 0 0	9 0	8 5	8 0
a-45	2 5 0	1 0 0	9 9	8 5	1 0 0
	1 2 5	1 0 0	9 0	5 0	9 0

Table 6 (continued)

Comp. No.	Dose of active ingredient (g/ha)	Growth-controlling degree (%)			
		EC	SJ	SP	OS
a-47	250	100	100	95	99
	125	100	85	60	70
a-48	250	100	95	90	95
	125	99	85	50	50
a-49	250	100	90	85	100
	125	100	90	50	100
a-51	250	100	99	85	90
	125	100	95	10	60
a-52	250	100	99	30	100
	125	100	90	30	95
a-53	125	100	95	90	80
	63	100	90	90	0
a-55	250	100	95	20	30
	125	100	50	0	30
a-58	250	100	85	50	40
	125	99	70	50	40
a-59	63	100	90	10	30
	31	100	50	-	30
a-60	250	100	100	70	95
	125	100	95	30	70
a-69	250	100	99	50	100
	125	100	85	50	35
a-72	125	90	85	20	40
	63	60	85	20	35
a-73	125	100	90	50	80
	63	100	70	50	30
a-76	125	100	90	30	0
	63	100	85	0	0
a-77	125	90	85	50	70
	63	70	50	30	40
a-78	250	100	85	85	80
	125	85	60	85	30
a-79	250	100	95	90	90
	125	100	85	50	40



Table 6 (continued)

Comp. No.	Dose of active ingredient (g/ha)	Growth-controlling degree (%)			
		EC	SJ	SP	OS
a-80	5 0 0	1 0 0	1 0 0	9 5	9 0
	2 5 0	9 9	1 0 0	9 5	9 0
a-81	5 0 0	9 5	9 9	9 0	3 0
	2 5 0	8 5	9 0	5 0	3 0
a-82	1 2 5	4 0	8 5	5 0	3 5
	6 3	2 0	1 0	0	2 0
a-83	2 5 0	9 9	9 9	9 0	6 0
	1 2 5	9 5	9 9	9 0	4 0
a-84	2 5 0	1 0 0	9 9	5 0	9 0
	1 2 5	1 0 0	9 5	5 0	5 0
a-85	2 5 0	1 0 0	8 5	5 0	2 0
	1 2 5	9 5	7 0	0	2 0
a-86	1 2 5	1 0	7 0	1 0	0
	6 3	1 0	6 0	1 0	0
a-87	2 5 0	1 0 0	9 0	3 0	4 0
	1 2 5	1 0 0	8 5	0	3 0
a-88	2 5 0	1 0 0	9 0	5 0	1 0 0
	1 2 5	1 0 0	9 0	2 0	9 5
a-89	2 5 0	1 0 0	9 9	9 0	9 9
	1 2 5	9 9	9 9	5 0	9 5
a-90	2 5 0	9 9	9 5	9 0	8 0
	1 2 5	9 9	9 0	8 5	7 0
a-91	5 0 0	9 9	9 9	9 0	3 5
	2 5 0	9 9	9 5	8 5	3 0
a-92	2 5 0	9 9	9 0	1 0	3 5
	1 2 5	8 5	8 5	0	1 0
a-93	2 5 0	1 0 0	1 0 0	—	4 0
	1 2 5	1 0 0	1 0 0	8 5	0
a-94	2 5 0	1 0 0	9 9	9 0	9 9
	1 2 5	1 0 0	9 5	1 0	9 9
a-95	2 5 0	1 0 0	1 0 0	—	—
	1 2 5	1 0 0	1 0 0	8 5	—
a-96	2 5 0	1 0 0	9 5	8 5	1 0 0
	1 2 5	1 0 0	9 0	1 0	9 9

Table 6 (continued)

Comp. No.	Dose of active ingredient (g/ha)	Growth-controlling degree (%)			
		EC	SJ	SP	OS
a-97	63 31	95 95	30 0	50 0	50 50
a-98	63 31	100 100	80 60	70 0	60 30
a-99	125 63	100 100	95 —	50 50	100 100
a-100	125 63	50 10	70 10	10 0	30 0
a-101	63 31	100 80	10 0	0 0	40 30
a-102	63 31	95 85	85 50	50 0	40 25
a-103	63 31	95 85	90 10	50 30	65 10
a-104	63 31	99 99	0 0	0 0	30 30
a-105	63 31	95 90	0 0	0 0	40 10
a-106	250 125	100 99	70 30	70 50	50 15
a-118	250 125	100 100	90 70	60 60	95 55
a-120	250 125	100 100	99 90	50 40	98 90
a-122	250 125	100 100	100 100	70 60	95 100
a-127	1000	20	60	70	—
a-131	250 125	100 100	90 85	— 90	80 50
a-132	250 125	100 100	85 50	— 50	— 30
a-135	250 125	100 70	20 0	0 0	10 0
a-138	125 63	100 80	20 —	0 0	5 15

Table 6 (continued)

Comp. No.	Dose of active ingredient (g/ha)	Growth-controlling degree (%)			
		EC	SJ	SP	OS
a-139	5 0 0	7 0	6 0	5 0	—
	2 5 0	7 0	2 0	2 0	—
a-140	2 5 0	1 0 0	9 0	3 0	9 8
	1 2 5	1 0 0	9 0	—	3 5
a-141	2 5 0	1 0 0	1 0 0	3 0	1 0 0
	1 2 5	1 0 0	1 0 0	0	9 5
a-143	5 0 0	1 0 0	1 0	3 0	—
	2 5 0	1 0 0	0	1 0	—
a-144	5 0 0	9 9	2 0	9 0	—
	2 5 0	7 0	1 0	8 5	—
a-145	2 5 0	1 0 0	6 0	5 0	0
	1 2 5	1 0 0	5 0	4 0	1 0
a-146	5 0 0	9 0	9 0	9 0	3 0
	2 5 0	6 0	7 0	7 0	0
a-148	1 2 5	1 0 0	7 0	0	0
	6 3	1 0 0	7 0	0	0
a-149	1 2 5	9 0	3 0	0	0
	6 3	8 5	1 0	0	0
a-150	2 5 0	1 0 0	5 0	6 0	0
	1 2 5	9 5	0	3 0	5
a-151	2 5 0	1 0 0	0	0	3 0
	1 2 5	1 0 0	0	—	5
a-152	2 5 0	1 0 0	6 0	0	1 0 0
	1 2 5	1 0 0	3 0	0	1 0 0
a-153	2 5 0	9 9	9 0	7 0	1 0
	1 2 5	1 0 0	3 0	3 0	2 0
a-154	2 5 0	1 0 0	2 0	4 0	6 0
	1 2 5	9 9	0	0	2 0
a-155	2 5 0	1 0 0	7 0	0	8 5
	1 2 5	1 0 0	6 0	0	5 0
a-157	2 5 0	9 0	9 5	7 0	0
	1 2 5	9 0	0	0	0
a-158	2 5 0	1 0 0	8 0	3 0	9 0
	1 2 5	1 0 0	5 0	6 0	9 0

Table 6 (continued)

Comp. No.	Dose of active ingredient (g/ha)	Growth-controlling degree (%)			
		EC	SJ	SP	OS
a-160	2 5 0 1 2 5	8 0 7 0	5 0 —	0 0	0 0
a-161	2 5 0 1 2 5	7 0 7 0	0 0	0 0	5 0
a-162	2 5 0 1 2 5	1 0 0 1 0 0	0 0	0 0	1 0 0
a-164	2 5 0 1 2 5	1 0 0 1 0 0	0 0	0 0	6 0 1 0
a-165	1 2 5 6 3	1 0 0 9 5	8 0 6 0	3 0 0	1 0 0 3 5
a-166	1 2 5 6 3	1 0 0 1 0 0	6 0 2 0	0 0	1 0 0 8 5
a-167	2 5 0 1 2 5	1 0 0 1 0 0	9 5 7 0	6 0 0	3 5 1 0
a-168	1 2 5 6 3	1 0 0 1 0 0	9 5 9 0	0 0	5 5 5 0
a-169	1 2 5 6 3	1 0 0 1 0 0	4 0 4 0	0 0	9 0 5 5
a-170	1 2 5 6 3	1 0 0 1 0 0	3 0 1 0	0 0	8 5 4 5
a-171	2 5 0	7 0	0	0	0
a-172	2 5 0	8 0	3 0	5 0	5 0
a-173	2 5 0 1 2 5	9 9 7 0	2 0 —	0 0	1 0 0 2 5
a-174	2 5 0 1 2 5	1 0 0 1 0 0	6 0 6 0	0 0	1 0 0 1 0 0
a-175	2 5 0 1 2 5	8 0 8 0	5 0 3 0	0 0	0 0
a-177	2 5 0 1 2 5	1 0 0 1 0 0	5 0 3 0	0 0	0 0
a-178	2 5 0 1 2 5	1 0 0 1 0 0	3 0 3 0	0 0	1 0 0
a-179	2 5 0 1 2 5	1 0 0 1 0 0	9 0 3 0	0 0	1 0 0 9 0

Table 6 (continued)

Comp. No.	Dose of active ingredient (g/ha)	Growth-controlling degree (%)			
		EC	SJ	SP	OS
a-180	2 5 0	1 0 0	3 0	0	8 0
	1 2 5	1 0 0	0	0	6 0
a-181	2 5 0	1 0 0	0	2 0	5 0
a-182	2 5 0	1 0 0	7 0	0	4 5
	1 2 5	1 0 0	0	0	1 5
a-184	2 5 0	8 5	0	0	0
a-185	2 5 0	7 0	0	0	0
a-187	2 5 0	7 0	8 0	2 0	1 5
	1 2 5	3 0	8 0	0	0
a-188	2 5 0	1 0 0	5 0	2 0	9 5
	1 2 5	1 0 0	5 0	0	8 0
a-189	2 5 0	1 0 0	5 0	2 0	5 5
	1 2 5	1 0 0	5 0	0	5 0
a-191	2 5 0	1 0 0	2 0	2 0	3 5
	1 2 5	9 5	0	0	0
a-194	2 5 0	9 0	3 0	0	1 0
a-200	2 5 0	1 0 0	9 0	0	7 0
	1 2 5	1 0 0	9 0	0	8 0
a-202	2 5 0	1 0 0	7 0	0	1 0
	1 2 5	8 5	2 0	0	1 0
a-203	2 5 0	1 0 0	1 0 0	9 5	1 0 0
	1 2 5	1 0 0	9 5	9 5	1 0 0
a-204	2 5 0	1 0 0	3 0	3 0	1 0
	1 2 5	9 9	0	0	5
a-205	2 5 0	8 5	0	0	5
a-206	2 5 0	1 0 0	0	6 0	1 0 0
	1 2 5	9 5	0	7 0	9 0
a-207	2 5 0	9 5	0	6 0	5 0
	1 2 5	8 0	0	8 0	2 5
a-208	2 5 0	1 0 0	3 0	7 0	8 0
	1 2 5	1 0 0	0	6 0	5 0
a-210	2 5 0	1 0 0	4 0	7 0	1 0 0
	1 2 5	1 0 0	7 0	6 0	8 5

124  
Table 6 (continued)

Comp. No.	Dose of active ingredient (g/ha)	Growth-controlling degree (%)			
		EC	SJ	SP	OS
a-213	2 5 0	1 0 0	5 0	0	9 5
	1 2 5	1 0 0	2 0	0	6 5
a-214	2 5 0	1 0 0	6 0	2 0	6 0
	1 2 5	9 9	6 0	0	1 5
a-216	2 5 0	1 0 0	8 0	0	9 5
	1 2 5	1 0 0	4 0	0	1 5
a-217	2 5 0	1 0 0	2 0	0	1 0 0
	1 2 5	9 5	0	0	5 0
a-218	2 5 0	1 0 0	7 0	2 0	1 0 0
	1 2 5	7 0	3 0	0	8 5
a-219	2 5 0	1 0 0	3 0	0	1 0 0
	1 2 5	1 0 0	0	0	7 0
a-230	6 3	1 0 0	0	0	—
	3 1	1 0 0	0	0	—
a-232	6 3	1 0 0	3 0	—	—
	3 1	1 0 0	0	0	—
a-233	6 3	7 0	0	0	—
	3 1	5 0	0	0	—
a-239	6 3	1 0 0	0	0	0
	3 1	8 0	0	0	0
a-249	2 5 0	1 0 0	0	0	9 5
	1 2 5	1 0 0	0	0	4 0
a-250	6 3	1 0 0	4 0	0	8 0
	3 1	1 0 0	0	0	4 5
a-253	6 3	1 0 0	0	0	3 5
	3 1	1 0 0	0	0	3 0
a-254	6 3	1 0 0	0	—	5
	3 1	1 0 0	0	0	0
a-256	6 3	1 0 0	0	0	0
	3 1	8 0	0	0	0
a-257	2 5 0	1 0 0	2 0	0	0
	1 2 5	1 0 0	0	0	0
a-258	2 5 0	6 0	0	2 0	5
	1 2 5	7 0	0	0	0

Table 6 (continued)

Comp. No.	Dose of active ingredient (g/ha)	Growth-controlling degree (%)			
		EC	SJ	SP	OS
a-259	2 5 0	1 0 0	0	0	1 0
	1 2 5	1 0 0	0	0	0
a-260	2 5 0	1 0 0	0	2 0	1 0
	1 2 5	8 5	0	0	0
a-261	2 5 0	1 0 0	0	4 0	5
	1 2 5	7 0	0	4 0	0
a-262	2 5 0	1 0 0	1 0	0	1 0
	1 2 5	1 0 0	0	0	0
a-263	2 5 0	1 0 0	0	0	1 0
	1 2 5	9 5	0	0	0
a-264	2 5 0	8 5	0	0	3 0
	1 2 5	7 0	0	0	1 5
a-271	6 3	1 0 0	0	0	—
	3 1	9 5	0	0	—
b-1	2 5 0	1 0 0	8 5	5 0	5 0
	1 2 5	1 0 0	5 0	5 0	2 0

Now, Formulation Examples of the present invention will be given. Compound Nos. in Formulation Examples correspond to Compound Nos. in Table 4a to 4b given hereinbefore.

5 FORMULATION EXAMPLE 1

- |    |   |                      |
|----|---|----------------------|
|    | (1) Compound No. a-12   | 75 parts by weight   |
|    | (2) Sodium N-methyl-N-oleoyl taurate<br>(Geropon T-77, tradename,<br>manufactured by Rhone-Poulenc) | 14.5 parts by weight |
|    | (3) NaCl  | 10 parts by weight   |
| 10 | (4) Dextrin   | 0.5 part by weight   |

The above components are placed in a high-speed mixing granulator, admixed with 20 wt% of water, granulated, and dried to form water-dispersible granules.

15 FORMULATION EXAMPLE 2

- |    |  |                    |
|----|--|--------------------|
|    | (1) Kaolin   | 78 parts by weight |
|    | (2) Condensate of sodium naphthalene<br>sulfonate and formalin (Laveline<br>FAN, tradename, manufactured by<br>Dai-ichi Kogyo Seiyaku Co., Ltd.)                   | 2 parts by weight  |
| 20 | (3) Sodium polyoxyethylene alkylaryl<br>ether sulfate-premix with white<br>carbon (Sorpul 5039, tradename,<br>manufactured by Toho Chemical<br>Industry Co., Ltd.) | 5 parts by weight  |
|    | (4) White carbon (Carplex, tradename,<br>manufactured by Shionogi Seiyaku<br>Co., Ltd.)  | 15 parts by weight |

25 The mixture of the above components (1) to (4) and Compound No. a-6 are mixed in a weight ratio of 9:1 to obtain a wettable powder.



127

## FORMULATION EXAMPLE 3

- |    |  |                    |
|----|--|--------------------|
|    | (1) Talc micropowder (Hi-Filler No. 10, tradename, manufactured by Matsumura Sangyo Co., Ltd.)   | 33 parts by weight |
| 5  | (2) Dialkyl sulfosuccinate-premixed with white carbon (Sorpul 5050, tradename, manufactured by Toho Chemical Industry Co., Ltd.)   | 3 parts by weight  |
|    | (3) A mixture of polyoxyethylene alkylaryl ether sulfate and a polyoxyethylene monomethyl ether carbonate, premixed with white carbon (Sorpul 5073, tradename, manufactured by Toho Chemical Industry Co., Ltd.) | 4 parts by weight  |
| 10 | (4) Compound No. a-42  | 60 parts by weight |

The above components (1) to (4) are mixed to obtain a wettable powder.

## FORMULATION EXAMPLE 4

- |    |  |                      |
|----|--|----------------------|
| 15 | (1) Compound No. a-27  | 4 parts by weight    |
|    | (2) Bentonite  | 30 parts by weight   |
|    | (3) Calcium carbonate  | 61.5 parts by weight |
|    | (4) Polycarboxylic acid type surfactant (Toxanon GR-31A, tradename, manufactured by Sanyo Chemical Industries Co., Ltd.) | 3parts by weight     |
| 20 | (5) Calcium lignin sulfonate   | 1.5 parts by weight  |

Pulverized component (1) and components (2) and (3) are preliminarily mixed, and then components (4) and (5) and water are mixed thereto. The mixture is extruded and  
25 granulated, followed by drying and size-adjusting to obtain granules.

## FORMULATION EXAMPLE 5


128

- (1) Compound No. a-22 30 parts by weight
- (2) A pulverized product of a mixture of kaolinite and sericite (Zieclite, tradename, manufactured by Zieclite Co., Ltd.) 60 parts by weight
- 5 (3) Alkyl naphthalene sulfonate (New Kalgen WG-1, tradename, manufactured by Takemoto Oils and Fats Co., Ltd.) 5 parts by weight
- (4) Polyoxyalkylene allyl phenyl ether sulfate (New Kalgen FS-7, tradename, manufactured by Takemoto Oils and Fats Co., Ltd.) 5 parts by weight
- 10 Components (1), (2) and (3) are mixed and passed through a pulverizer, and then component (4) is added thereto. The mixture is kneaded and then extruded and granulated, followed by drying and size-adjusting to obtain water-dispersible granules.
- 15 FORMULATION EXAMPLE 6
- (1) Compound No. a-13 28 parts by weight
- (2) Triethanolamine salts of oxyethylated polyarylphenol phosphate (Soprophor FL, tradename, manufactured by Rhone-Poulenc) 2 parts by weight
- 20 (3) A mixture of polyoxyethylene styryl phenyl ether and alkyl aryl sulfonate (Sorpul 355, tradename, manufactured by Toho Chemical Industry Co., Ltd.) 1 part by weight
- (4) Isoparaffin hydrocarbon (IP solvent 1620, tradename, manufactured by Idemitsu Petrochemical Co., Ltd.) 32 parts by weight
- 25 (5) Ethylene glycol 6 parts by weight
- (6) Water 31 parts by weight

The above components (1) to (6) are mixed and pulverized by a wet-grinding machine (Dyno-mill) to obtain a water-based suspension concentrate.

CLAIMS

1. A pyrazole compound of the formula (I) or its salt:




wherein  $R_1$  is an alkyl group,  $R_2$  is a hydrogen atom, a  
 10 methyl group,  $-A-R_3$ , a phenyl group which may be  
 substituted, a pyridyl group which may be substituted, or  
 an allyl group which is substituted by a phenyl group, A  
 is  $-SO_2-$ ,  $-CO-$ ,  $-CH(R_6)-$  or  $-CH(R_7)CO-$ ,  $R_3$  is an alkyl  
 group which may be substituted, an alkenyl group which  
 15 may be substituted, an alkynyl group which may be  
 substituted, an alkoxy group which may be substituted, a  
 cyano group, a dialkylamino group or a phenyl group which  
 may be substituted, each of  $R_6$  and  $R_7$  is a hydrogen atom  
 or an alkyl group, X is a hydrogen atom, a halogen atom,  
 20 an alkyl group, a haloalkyl group, an alkoxy group, an  
 alkylthio group, an alkylsulfinyl group, an alkylsulfonyl  
 group, a nitro group, an alkoxycarbonyl group,  
 $-SO_2N(R_8)R_9$ ,  $-N(R_{10})SO_2R_{11}$ ,  $-CH_2S(O)qR_{12}$  or  $-OSO_2R_{13}$ , each  
 of  $R_8$ ,  $R_9$ ,  $R_{10}$ ,  $R_{11}$ ,  $R_{12}$  and  $R_{13}$  is an alkyl group, Z is  
 25 an alkyl group, l is an integer of from 0 to 5, n is an  
 integer of from 1 to 5, and q is an integer of from 0 to  
 2, provided that when l is at least 2, a plurality of Z

may be the same or different, and when n is at least 2, a plurality of X may be the same or different.

2. The pyrazole compound or its salt according to Claim 1, wherein the formula (I) is represented by the formula

5 (I'):



(I')

10 wherein  $R_1$  is an alkyl group,  $R_2$  is a hydrogen atom or  $-A-R_3$ , A is  $-SO_2-$ ,  $-CO-$ ,  $-CH_2-$  or  $-CH_2CO-$ ,  $R_3$  is an alkyl group which may be substituted, an alkenyl group which may be substituted, an alkynyl group which may be substituted, a cyano group or a phenyl group which may be substituted, each of  $X^1$ ,  $X^2$  and  $X^3$  is a hydrogen atom, a halogen atom, an alkyl group, a haloalkyl group, an alkoxy group, an alkylthio group, an alkylsulfinyl group, an alkylsulfonyl group, a nitro group,  $-SO_2N(R_8)R_9$ ,  $-N(R_{10})SO_2R_{11}$ ,  $-CH_2S(O)qR_{12}$  or  $-OSO_2R_{13}$ , each of  $R_8$ ,  $R_9$ ,  $R_{10}$ ,  $R_{11}$ ,  $R_{12}$  and  $R_{13}$  is an alkyl group, and q is an integer of from 0 to 2.


25 3. The pyrazole compound or its salt according to Claim 2, wherein A is  $-SO_2-$ ,  $-CH_2-$  or  $-CH_2CO-$ , each of  $X^1$ ,  $X^2$  and  $X^3$  is a hydrogen atom, a halogen atom, an alkyl group, a haloalkyl group, an alkoxy group, an alkylthio

group, an alkylsulfinyl group, an alkylsulfonyl group or a nitro group.

4. The pyrazole compound or its salt according to Claim 3, wherein  $X^1$  is an alkylthio group, an alkylsulfinyl group or an alkylsulfonyl group, and each of  $X^2$  and  $X^3$  is a hydrogen atom, a halogen atom, an alkyl group, a haloalkyl group or a nitro group.

5. A process for producing a pyrazole compound of the formula (I-1) or its salt:

10



(I-1)


15

wherein  $R_1$  is an alkyl group,  $Z$  is an alkyl group,  $l$  is an integer of from 0 to 5, provided that when  $l$  is at least 2, a plurality of  $Z$  may be the same or different,  $X$  is a hydrogen atom, a halogen atom, an alkyl group, a haloalkyl group, an alkoxy group, an alkylthio group, an alkylsulfinyl group, an alkylsulfonyl group, a nitro group, an alkoxycarbonyl group,  $-SO_2N(R_8)R_9$ ,  $-N(R_{10})SO_2R_{11}$ ,  $-CH_2S(O)qR_{12}$  or  $-OSO_2R_{13}$ , each of  $R_8$ ,  $R_9$ ,  $R_{10}$ ,  $R_{11}$ ,  $R_{12}$  and  $R_{13}$  is an alkyl group,  $q$  is an integer of from 0 to 2, and  $n$  is an integer of from 1 to 5, provided that when  $n$  is at least 2, a plurality of  $X$  may be the same or different, which comprises reacting a


25

133


compound of the formula (II):



wherein  $R_1$ ,  $Z$  and  $l$  are as defined above, with a compound of the formula (III):



wherein  $X$  and  $n$  are as defined above, and  $Y$  is a halogen atom, to obtain a compound of the formula (IV):




20

wherein  $R_1$ ,  $X$ ,  $Z$ ,  $l$  and  $n$  are as defined above, and subjecting the compound of the formula (IV) to a rearrangement reaction.

6. A process for producing a pyrazole compound of the  
25 formula (I-1') or its salt:


134



5

wherein  $R_1$  is an alkyl group,  $Z$  is an alkyl group,  $l$  is an integer of from 0 to 5, provided that when  $l$  is at least 2, a plurality of  $Z$  may be the same or different,  $X'$  is a hydrogen atom, a halogen atom, an alkyl group, a haloalkyl group, an alkoxy group, an alkylthio group, an alkylsulfinyl group or an alkylsulfonyl group, and  $n$  is an integer of from 1 to 5, provided that when  $n$  is at least 2, a plurality of  $X'$  may be the same or different, which comprises reacting a compound of the formula (II):

15



20 wherein  $R_1$ ,  $Z$  and  $l$  are as defined above, with a compound of the formula (V):




25 wherein  $X'$  and  $n$  are as defined above, and carbon tetrachloride, followed by a hydrolytic reaction.

7. A process for producing a pyrazole compound of the




135

formula (I-1) or its salt:



wherein  $R_1$  is an alkyl group,  $Z$  is an alkyl group,  $l$  is an integer of from 0 to 5, provided that when  $l$  is at least 2, a plurality of  $Z$  may be the same or different,  $X$  is a hydrogen atom, a halogen atom, an alkyl group, a haloalkyl group, an alkoxy group, an alkylthio group, an alkylsulfinyl group, an alkylsulfonyl group, a nitro group, an alkoxy carbonyl group,  $-\text{SO}_2\text{N}(\text{R}_8)\text{R}_9$ ,  $-\text{N}(\text{R}_{10})\text{SO}_2\text{R}_{11}$ ,  $-\text{CH}_2\text{S}(\text{O})\text{qR}_{12}$  or  $-\text{OSO}_2\text{R}_{13}$ , each of  $\text{R}_8$ ,  $\text{R}_9$ ,  $\text{R}_{10}$ ,  $\text{R}_{11}$ ,  $\text{R}_{12}$  and  $\text{R}_{13}$  is an alkyl group,  $q$  is an integer of from 0 to 2, and  $n$  is an integer of from 1 to 5, provided that when  $n$  is at least 2, a plurality of  $X$  may be the same or different, which comprises reacting a compound of the formula (II):

20



wherein  $R_1$ ,  $Z$  and  $l$  are as defined above, with a compound

136


of the formula (VI):



5

wherein X and n are as defined above.

8. A process for producing a pyrazole compound of the formula (I-1) or its salt:




15 wherein  $\text{R}_1$  is an alkyl group, Z is an alkyl group, l is an integer of from 0 to 5, provided that when l is at least 2, a plurality of Z may be the same or different, X is a hydrogen atom, a halogen atom, an alkyl group, a haloalkyl group, an alkoxy group, an alkylthio group, an alkylsulfinyl group, an alkylsulfonyl group, a nitro group, an alkoxycarbonyl group,  $-\text{SO}_2\text{N}(\text{R}_8)\text{R}_9$ ,  $-\text{N}(\text{R}_{10})\text{SO}_2\text{R}_{11}$ ,  $-\text{CH}_2\text{S}(\text{O})_q\text{R}_{12}$  or  $-\text{OSO}_2\text{R}_{13}$ , each of  $\text{R}_8$ ,  $\text{R}_9$ ,  $\text{R}_{10}$ ,  $\text{R}_{11}$ ,  $\text{R}_{12}$  and  $\text{R}_{13}$  is an alkyl group, q is an integer of from 0 to 2, and n is an integer of from 1 to 5, provided that when n is at least 2, a plurality of X may be the same or different, which comprises reacting a compound of the formula (II):


20

25

137




wherein  $R_1$ , Z and l are as defined above, with a compound of the formula (X):



wherein X and n are as defined above, and T is a chlorine atom, a bromine atom or an iodine atom, and carbon monoxide.


- 15 9. A process for producing a pyrazole compound of the formula (I-2) or its salt:



wherein  $R_1$  is an alkyl group,  $R_2'$  is a methyl group,  $-A-R_3$ , a phenyl group which may be substituted, a pyridyl group which may be substituted or an allyl group which is substituted by a phenyl group, A is  $-SO_2-$ ,  $-CO-$ ,  $-CH(R_6)-$  or  $-CH(R_7)CO-$ ,  $R_3$  is an alkyl group which may be

25

substituted, an alkenyl group which may be substituted,  
 an alkynyl group which may be substituted, an alkoxy  
 group which may be substituted, a cyano group, a  
 dialkylamino group or a phenyl group which may be  
 5 substituted, each of  $R_6$  and  $R_7$  is a hydrogen atom or an  
 alkyl group,  $Z$  is an alkyl group,  $l$  is an integer of from  
 0 to 5, provided that when  $l$  is at least 2, a plurality  
 of  $Z$  may be the same or different,  $X$  is a hydrogen atom,  
 a halogen atom, an alkyl group, a haloalkyl group, an  
 10 alkoxy group, an alkylthio group, an alkylsulfinyl group,  
 an alkylsulfonyl group, a nitro group, an alkoxycarbonyl  
 group,  $-\text{SO}_2\text{N}(\text{R}_8)\text{R}_9$ ,  $-\text{N}(\text{R}_{10})\text{SO}_2\text{R}_{11}$ ,  $-\text{CH}_2\text{S}(\text{O})_q\text{R}_{12}$  or  
 $-\text{OSO}_2\text{R}_{13}$ , each of  $R_8$ ,  $R_9$ ,  $\text{R}_{10}$ ,  $\text{R}_{11}$ ,  $\text{R}_{12}$  and  $\text{R}_{13}$  is an alkyl  
 group and  $n$  is an integer of from 1 to 5, provided that  
 15 when  $n$  is at least 2, a plurality of  $X$  may be the same or  
 different,  $q$  is an integer of from 0 to 2, which  
 comprises reacting a compound of the formula (I-1):




where  $\text{R}_1$ ,  $X$ ,  $Z$ ,  $n$  and  $l$  are as defined above, with a  
 25 compound of the formula (VII):



wherein  $\text{R}_2'$  is as defined above, and  $\text{Y}$  is a halogen atom.


10. A process for producing a pyrazole compound of the formula (I-4) or its salt:



wherein each of  $R_1$  and  $R_5$  is an alkyl group,  $Z$  is an alkyl group,  $l$  is an integer of from 0 to 5, provided that when  $l$  is at least 2, a plurality of  $Z$  may be the same or different,  $X''$  is a hydrogen atom, a halogen atom, an alkyl group, a haloalkyl group, an alkoxy group, an alkylsulfinyl group, an alkylsulfonyl group, a nitro group, an alkoxycarbonyl group,  $-\text{SO}_2\text{N}(R_8)R_9$ ,  $-\text{N}(R_{10})\text{SO}_2R_{11}$ ,  $-\text{CH}_2\text{S}(\text{O})_qR_{12}$  or  $-\text{OSO}_2R_{13}$ , each of  $R_8$ ,  $R_9$ ,  $R_{10}$ ,  $R_{11}$ ,  $R_{12}$  and  $R_{13}$  is an alkyl group,  $q'$  is 1 or 2,  $m$  is 1 or 2, and  $n$  is an integer of from 1 to 5, provided that when  $n$  is at least 3, a plurality of  $X''$  may be the same or different, which comprises oxidizing a compound of the formula (VI-1):


15

20



140

where  $R_1$ ,  $R_5$ ,  $Z$ ,  $l$  and  $n$  are as defined above and  $X$  is a hydrogen atom, a halogen atom, an alkyl group, a haloalkyl group, an alkoxy group, an alkylthio group, an alkylsulfinyl group, an alkylsulfonyl group, a nitro group, an alkoxy carbonyl group,  $-SO_2N(R_8)R_9$ ,  $-N(R_{10})SO_2R_{11}$ ,  $-CH_2S(O)_qR_{12}$  or  $-OSO_2R_{13}$ , each of  $R_8$ ,  $R_9$ ,  $R_{10}$ ,  $R_{11}$ ,  $R_{12}$  and  $R_{13}$  is an alkyl group, provided that when  $n$  is at least 3, a plurality of  $X$  may be the same or different to obtain a compound of the formula (VI-2):




15

wherein  $R_1$ ,  $R_5$ ,  $Z$ ,  $X''$ ,  $l$ ,  $m$  and  $n$  are as defined above, and subjecting the compound of the formula (VI-2) to a rearrangement reaction.


11. A process for producing a pyrazole compound of the formula (I-7) or its salt:

20



141

wherein each of  $R_1$  and  $R_5$  is an alkyl group,  $R_2'$  is a methyl group,  $-A-R_3$ , a phenyl group which may be substituted, a pyridyl group which may be substituted or an allyl group which is substituted by a phenyl group, A is  $-SO_2-$ ,  $-CO-$ ,  $-CH(R_6)-$  or  $-CH(R_7)CO-$ ,  $R_3$  is an alkyl group which may be substituted, an alkenyl group which may be substituted, an alkynyl group which may be substituted, an alkoxy group which may be substituted, a cyano group, a dialkylamino group or a phenyl group which may be substituted, each of  $R_6$  and  $R_7$  is a hydrogen atom or an alkyl group, Z is an alkyl group, l is an integer of from 0 to 5, provided that when l is at least 2, a plurality of Z may be the same or different, m is 1 or 2,  $X''$  is a hydrogen atom, a halogen atom, an alkyl group, a haloalkyl group, an alkoxy group, an alkylsulfinyl group, an alkylsulfonyl group, a nitro group, an alkoxycarbonyl group,  $-SO_2N(R_8)R_9$ ,  $-N(R_{10})SO_2R_{11}$ ,  $-CH_2S(O)_{q'}R_{12}$  or  $-OSO_2R_{13}$ , each of  $R_8$ ,  $R_9$ ,  $R_{10}$ ,  $R_{11}$ ,  $R_{12}$  and  $R_{13}$  is an alkyl group,  $q'$  is 1 or 2, and n is an integer of from 1 to 5, provided that when n is at least 3, a plurality of  $X''$  may be the same or different, which comprises oxidizing a compound of the formula (I-6):




- where  $R_1$ ,  $R_2'$ ,  $R_5$ ,  $Z$ ,  $l$  and  $n$  are as defined above, and  $X$  is a hydrogen atom, a halogen atom, an alkyl group, a haloalkyl group, an alkoxy group, an alkylthio group, an alkylsulfinyl group, an alkylsulfonyl group, a nitro group, an alkoxycarbonyl group,  $-\text{SO}_2\text{N}(R_8)R_9$ ,  $-\text{N}(R_{10})\text{SO}_2R_{11}$ ,  $-\text{CH}_2\text{S}(\text{O})_qR_{12}$  or  $-\text{OSO}_2R_{13}$ , each of  $R_8$ ,  $R_9$ ,  $R_{10}$ ,  $R_{11}$ ,  $R_{12}$  and  $R_{13}$  is an alkyl group,  $q$  is an integer of from 0 to 2, provided that when  $n$  is at least 3, a plurality of  $X$  may be the same or different.
- 5
- 10 12. A herbicide containing the pyrazole compound or its salt as defined in Claim 1, as an active ingredient.
13. A method for controlling noxious weeds, which comprises applying an effective amount of the pyrazole compound or its salt as defined in Claim 1.
- 15 14. A method for controlling noxious weeds, which comprises applying an effective amount of the pyrazole compound or its salt as defined in Claim 1 to an upland field.
- 20 15. A method for controlling noxious weeds, which comprises applying an effective amount of the pyrazole compound or its salt as defined in Claim 1 to a corn field.
- 25 16. A method for controlling noxious weeds, which comprises applying an effective amount of the pyrazole compound or its salt as defined in Claim 1 to a wheat field.
17. A method for controlling noxious weeds, which



comprises applying an effective amount of the pyrazole compound or its salt as defined in Claim 1 to a paddy field.


18. A mixed herbicidal composition comprising at least one member selected from the pyrazole compound or its salt as defined in Claim 1 and at least one member selected from active ingredient compounds of other herbicides.

19. A compound of the formula (II):



15 wherein  $R_1$  is an alkyl group, Z is an alkyl group and l is an integer of from 0 to 5, provided that when l is at least 2, a plurality of Z may be the same or different.

20. The compound of the formula (IV):



wherein  $R_1$  is an alkyl group, X is a hydrogen atom, a

halogen atom, an alkyl group, a haloalkyl group, an alkoxy group, an alkylthio group, an alkylsulfinyl group, an alkylsulfonyl group, a nitro group, an alkoxycarbonyl group,  $-\text{SO}_2\text{N}(\text{R}_8)\text{R}_9$ ,  $-\text{N}(\text{R}_{10})\text{SO}_2\text{R}_{11}$ ,  $-\text{CH}_2\text{S}(\text{O})_q\text{R}_{12}$  or  
5  $-\text{OSO}_2\text{R}_{13}$ , each of  $\text{R}_8$ ,  $\text{R}_9$ ,  $\text{R}_{10}$ ,  $\text{R}_{11}$ ,  $\text{R}_{12}$  and  $\text{R}_{13}$  is an alkyl group,  $\text{Z}$  is an alkyl group,  $l$  is an integer of from 0 to 5,  $n$  is an integer of from 1 to 5, and  $q$  is an integer of from 0 to 2, provided that when  $l$  is at least 2, a plurality of  $\text{Z}$  may be the same or different, and when  $n$   
10 is at least 2, a plurality of  $\text{X}$  may be the same or different.

# INTERNATIONAL SEARCH REPORT

<b>A. CLASSIFICATION OF SUBJECT MATTER</b> IPC 6 C07D231/20 C07D401/12 A01N43/56		International Application No PCT/JP 97/01457
According to International Patent Classification (IPC) or to both national classification and IPC		
<b>B. FIELDS SEARCHED</b> Minimum documentation searched (classification system followed by classification symbols) IPC 6 C07D		
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched		
Electronic data base consulted during the international search (name of data base and, where practical, search terms used)		
<b>C. DOCUMENTS CONSIDERED TO BE RELEVANT</b>		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	CHEMICAL ABSTRACTS, vol. 114, no. 7, 18 February 1991 Columbus, Ohio, US; abstract no. 62091m, T. AONO ET AL.: "Preparation of pyrazolones and peroxy lipid formation inhibitors, lipoxygenase inhibitors, and collagenase inhibitors containing them." page 690; column 2; XP002037186 see abstract; and Chemical Abstracts, CHEMICAL SUBSTANCES, 12th Collective Index, vol. 106-115, 1987-1991, page 78553CS, the compounds with the RN: [131645-10-8], [131645-56-2], and [131645-19-7] & JP 02 229 168 A (TAKEDA CHEMICAL INDUSTRIES, LTD.) 11 September 1990 --- -/--	19
<input checked="" type="checkbox"/> Further documents are listed in the continuation of box C.		<input checked="" type="checkbox"/> Patent family members are listed in annex.
* Special categories of cited documents : "A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier document but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art "Z" document member of the same patent family		
Date of the actual completion of the international search 7 August 1997		Date of mailing of the international search report 26.08.97
Name and mailing address of the ISA European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Tx. 31 651 epo nl, Fax (+31-70) 340-3016		Authorized officer Fink, D

# INTERNATIONAL SEARCH REPORT

International Application No  
PCT/JP 97/01457

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	US 4 063 925 A (KONOTSUNE TAKUO ET AL) 20 December 1977 see column 1, line 7 - column 2, line 10 see column 19; example 3 ---	1-4, 12-18
Y	EP 0 638 555 A (NISSAN CHEMICAL IND LTD) 15 February 1995 cited in the application see page 149 - page 150; claims ---	1-4, 12-18
A	GB 2 002 375 A (ISHIHARA MINING & CHEMICAL CO) 21 February 1979 cited in the application see the whole document -----	1-20

# INTERNATIONAL SEARCH REPORT

Information on patent family members

Inter. Appl. Application No

PCT/JP 97/01457

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 4063925 A	20-12-77	JP 1001829 C	19-06-80
		JP 50126830 A	06-10-75
		JP 54036648 B	10-11-79
		AU 7934175 A	23-09-76
		CA 1077944 A	20-05-80
		CA 1069329 A	08-01-80
		CA 1075030 A	08-04-80
		DE 2513750 A	09-10-75
		FR 2274219 A	09-01-76
		GB 1463473 A	02-02-77
		US 4414392 A	08-11-83
		US 4301293 A	17-11-81
		US 4508910 A	02-04-85
		US 4687858 A	18-08-87
		US 4146726 A	27-03-79
		US 4261729 A	14-04-81
EP 0638555 A	15-02-95	AU 673505 B	14-11-96
		AU 2933092 A	15-06-93
		BR 9206728 A	21-11-95
		US 5545608 A	13-08-96
		CA 2123499 A	27-05-93
		CN 1072177 A	19-05-93
		WO 9310099 A	27-05-93
		JP 6184115 A	05-07-94
		CN 1077450 A	20-10-93
GB 2002375 A	21-02-79	JP 1252061 C	26-02-85
		JP 54041872 A	03-04-79
		JP 56028885 B	04-07-81
		JP 1305551 C	28-02-86
		JP 54070269 A	05-06-79
		JP 60029388 B	10-07-85
		US 4230481 A	28-10-80